

FOREST HEALTH IN NORTH AMERICA: SOME PERSPECTIVES ON ACTUAL AND POTENTIAL ROLES OF CLIMATE AND AIR POLLUTION

S. McLAUGHLIN¹ and K. PERCY²

¹Environmental Sciences Division, Oak Ridge National Laboratory, PO Box 2008, MS 6422, Oak Ridge, TN, 37830, USA; ²Natural Resources Canada, Canadian Forest Service, Atlantic Forestry Centre, Fredericton, N.B., E3B 5P7, Canada

(Received 25 September 1998; accepted 25 February 1999)

Abstract. The perceived health of forest ecosystems over large temporal and spatial scales can be strongly influenced by the frames of reference chosen to evaluate both forest condition and the functional integrity of sustaining forest processes. North American forests are diverse in range, species composition, past disturbance history, and current management practices. Therefore the implications of changes in environmental stress from atmospheric pollution and/or global climate change on health of these forests will vary widely across the landscape. Forest health surveys that focus on the average forest condition may do a credible job of representing the near-term trends in economic value while failing to detect fundamental changes in the processes by which these values are sustained over the longer term. Indications of increased levels of environmental stress on forest growth and nutrient cycles are currently apparent in several forest types in North America. Measurements of forest ecophysiological responses to air pollutants in integrated case studies with four forest types (southern pine, western pine, high elevation red spruce, and northeastern hardwoods) indicate that ambient levels of ozone and/or acidic deposition can alter basic processes of water, carbon, and nutrient allocation by forest trees. These changes then provide a mechanistic basis for pollutant stress to enhance a wider range of natural stresses that also affect and are affected by these resources. Future climatic changes may ameliorate (+ CO₂) or exacerbate (+ temperature, + UV-B) these effects. Current projections of forest responses to global climate change do not consider important physiological changes induced by air pollutants that may amplify climatic stresses. These include reduced rooting mass, depth, and function, increased respiration, and reduced water use efficiency. Monitoring and understanding the relative roles of natural and anthropogenic stress in influencing future forest health will require programs that are structured to evaluate responses at appropriate frequencies across gradients in both forest resources and the stresses that influence them. Such programs must also be accompanied by supplemental process-oriented and pattern-oriented investigations that more thoroughly test cause and effect relationships among stresses and responses of both forests and the biogeochemical cycles that sustain them.

Key Words: air quality, physiology, growth, biotic and abiotic interactions

1. Introduction

The forests of North America are immense in both their geographical range and ecological diversity. The 326 M ha of forested land in the USA (Powell *et al.*, 1992) and 417 M ha in Canada (NRCAN, 1998) are also a highly valuable economic resource for which maintenance of long term productivity is a very high priority. Both actual and perceived potential responses of North American forests to atmospheric pollution during recent decades have figured strongly in policy decisions on air quality regulation that have had significant economic and ecological implications for the region. In the next century, if current increases in atmospheric CO₂ continue and projected changes in both temperature and atmospheric deposition of nitrogen (N) occur, changes in forest productivity and function will be of interest not only in terms of shifts in patterns of production, but also from the perspective of the role of forests as sinks or sources for greenhouse gases (Norby, 1998). IPCC (1998) predictions for the effects of Global Climate Change on forest health include both increased growth and range of some forests, but also an increased frequency of declining health in others in response to increased biotic and abiotic stresses associated

The authors wish to express their sincere appreciation to Dr. R. Vet, Atmospheric Environment Service, Environment Canada, Downsview, Ontario, and to Dr. T. Dann, Environment Protection Service, Environment Canada, Ottawa, Ontario, for calculation and mapping of SO₄, NO₃H ion depositions and cumulative O₃ sums.

Water, Air, and Soil Pollution 116: 151-197, 1999.

© 1999 Kluwer Academic Publishers. Printed in the Netherlands.

with climatic warming. Evaluating forest health and forest biogeochemical functions in this more complex future environment will increasingly challenge forest scientists to measure and understand the basic processes underlying forest growth and longer term ecological health.

Perspectives on forest health and current risks posed by chemical and physical climate change differ widely within the forest community (Jenkins, 1997). These differences often stem from varying perspectives of forest values and times scales over which change is evaluated. Kolb *et al.* (1994) differentiated between the utilitarian view of forest health, which stresses timber production as the primary endpoint and the ecosystem perspective, which views forest health in terms of measures of longer term forest function. The latter include properties such as resilience, diversity, and the flow of carbon, water, and nutrient resources required for tree resistance to natural stresses and maintenance of biogeochemical cycles. Assessment of forest health, in fact, has many dimensions (Innes, 1993), and at the time of our writing, there exists no universally-accepted definition of a healthy forest. O'Laughlin *et al.* (1994) have defined forest health as "...a condition of forest ecosystems that sustains their complexity while providing for human needs". This definition stresses the utilitarian endpoint, which is defined in anthropocentric terms. We focus here on a more fundamental definition - A capacity to supply and allocate water, nutrients, and energy in ways that increase or maintain productivity while maintaining resistance to biotic and abiotic stresses. We, therefore, define less healthy forests as ones in which trees lose productive capacity and/or become more sensitive to environmental stresses. We should note here that unhealthy trees and stands occur naturally as a part of successional processes by which a balance between forest production, site resources, and climate are attained. In this capacity, for example, grazing insects can serve an important role in establishing a balance in nutrient flow to foliage from nutrient deficient soils (Matson and Addy, 1975). The patterns by which such changes occur however are typically spatially and temporally heterogeneous as are the localized biotic and abiotic factors that influence them.

Ecosystem response to environmental stresses, including air pollutants, is a complex, hierarchical process occurring over time scales ranging from minutes (leaf) to decades (stand). However, air pollution stresses are somewhat distinct from many other forms of environmental stress because of their regional patterns and temporal consistency with changes in air quality. Stress responses begin with sensitive individuals proceeding from branch, to tree, and then stand and ecosystem levels (Hinckley *et al.*, 1992). Underlying these response are process-level changes in uptake and allocation of carbon, water, and nutrient resources. We focus here on linkages between process level changes and ecosystem responses because this approach offers the best chance to understand the individual and interactive effects that will ultimately determine the utilitarian values of forest systems.

This paper summarizes recent advances in understanding of process level effects of regional air pollution on four forest types for which recent syntheses are available. These include the southeastern pine forest, eastern spruce/fir ecosystem, northeastern hardwoods, and western pine forests in Southern California (Figure 1). Our objective is to evaluate how increased understanding of forest process effects can be integrated into regional monitoring and analysis of forest health in relation to both air pollution and its interactions with other biotic and abiotic stresses that affect forests. We have emphasized primarily ozone and acidic deposition (N and S), the pollutants with the greatest potential regional impact on forests in the U.S. and Canada. Measured and potential impacts have been evaluated from the following perspectives:




Figure 1. Multidisciplinary research from case studies in southwestern pine, southeastern pine, eastern spruce-fir, and northeastern hardwood forest types was examined to evaluate process level responses to ambient levels of air pollutants in four regional forest types.

- (1). Monitoring air quality and forest health in North America.
- (2). Evidence of changes in allocation patterns of carbon, water, and nutrients related to ozone or acidic deposition within the four "case-study" forests.
- (3). Implications of process-level changes on whole-tree function and biogeochemical processes in current and future climate regimes.
- (4). Recommendations for incorporating process-based ecosystem perspectives into regional monitoring of forest health.

2. Regional Air Quality in North America

Recent syntheses in both the United States (NAPAP, 1998) and Canada (EC, 1997) demonstrate clear qualitative relationships between spatial patterns of acidic gas emissions and measured air concentrations and wet deposition patterns that were already evident by the 1990 assessments (RMCC, 1990). Here, we concentrate on acidic deposition and O_3 as they affect our forests regionally, while recognizing that their co-occurrence and interactions with other abiotic (e.g., climate extremes, UV-B, toxics) and biotic stressors ultimately determines the direction and magnitude of forest health responses.

2.1. ACIDIC DEPOSITION

Emissions of SO_2 in eastern North America have declined from over 20 M t in 1980 to under 17 M t in 1993 (EC, 1997). As a consequence, US air concentrations of SO_2 have decreased 37% between 1985 and 1995, particularly in the northeast (NAPAP, 1998). Particulate SO_4 concentration reductions have been widespread, except at several mid- to high-elevation, forested areas (NAPAP, 1998). US emissions of anthropogenic NO_x decreased 6.5% from 23.3 M t in 1980 to 21.8 M t in 1995 (NAPAP, 1998). Trends in N species contributing to acidic deposition (HNO_3 , NO_3) were more variable than for SO_2 . While decreasing HNO_3 was reported in the northeastern US (NAPAP, 1998), Canadian data showed an increasing frequency of higher concentration HNO_3 events (EC, 1997).

Trends in distribution patterns for wet deposition of SO_4 and NO_3 are shown in Figures 2 and 3. It should be noted that additional inputs of dry deposition may add 8-37% more S and 15-65% more N to these numbers depending on the region (EC, 1997). Contrasts in wet SO_4 and NO_3 deposition patterns reveal strong reductions for S in the area enclosed by the 10-30 $kg\ ha^{-1}\ yr^{-1}$ isopleths. Also evident was the disappearance of the 30 $kg\ ha^{-1}\ yr^{-1}$ area in Canada, and its considerable reduction in the US (Figures 2a,b) (EC, 1997). In contrast, NO_3 wet deposition did not change dramatically. Areas receiving 5-25 $kg\ NO_3\ ha^{-1}\ yr^{-1}$ expanded marginally (Figures 3 a,b) (EC, 1997). When integrated, areas in Canada within the 20 $kg\ ha^{-1}\ yr^{-1}$ wet SO_4 isopleth decreased 44% from 629,000 ha in 1980-83 to 340,000 ha in 1990-93 (EC, 1997).

Hall *et al.* (1997) recommended a critical threshold of 12 $kg\ SO_4\ ha^{-1}\ yr^{-1}$ for wet deposition to poorly buffered Canadian soils. In the context of forest/lake hydrological process linkages, current wet SO_4 deposition exceeds critical loads at all eastern Canadian lake clusters by 7-12 $kg\ ha^{-1}\ yr^{-1}$ and is expected to exceed them by about 6 to 10 $kg\ ha^{-1}\ yr^{-1}$ after all emission controls required by the Canada/US Air Quality Agreement are fully implemented (Jeffries, 1997).




Figure 2. Four-year mean wet deposition patterns of SO_4^{2-} ($\text{kg ha}^{-1}\text{yr}^{-1}$) in eastern North America for (a) 1980-1983 and (b) 1990-1993. Reproduced from EC (1997).

Areas in Canada within the $15 \text{ kg wet NO}_3 \text{ ha}^{-1}\text{yr}^{-1}$ isopleth increased 13% from 373,000 ha in 1980-83 to 420,000 ha in 1990-93. Areas receiving at least $0.4 \text{ kg H}^+ \text{ ha}^{-1}\text{yr}^{-1}$ decreased along the east coast and in southern Ontario, while the area receiving at least $0.2 \text{ kg ha}^{-1}\text{yr}^{-1}$ expanded to the northeast (Newfoundland) and east (Nova Scotia) (Figures 4a,b) (EC, 1997). NO_3 and NH_4 concentrations showed high variability, and deposition of NH_4 , generally increased throughout the US (NAPAP, 1998). Acidification potential remains high over large forest areas due to increasing NO_3 and decreasing Ca, Mg, K, Na deposition (Likens *et al.*, 1996).

Episodes have a large impact upon temporal changes in annual deposition amounts (Brook, 1995) and may increase annual mean SO_4 and H^+ deposition by $\leq 20\%$ (Beattie and Whelpdale, 1989). In this context, precipitation amount and type are critical in pollutant deposition to forests, particularly those growing at mid to high elevation. Most mountain forests receive more ($>50\%$) precipitation than those at lower elevation; coastal forests are immersed in fog for up to 30% of the growing season. Mountain cloud/fog and coastal fog have been found to have extremely low mean annual weighted pH's, consistently 10-50X more acidic than precipitation (Kimball *et al.*, 1988), with single events recorded as low as




Figure 3. Four-year mean wet deposition patterns of NO₃⁻ (kg ha⁻¹yr⁻¹) in eastern North America for (a) 1980-1983 and (b) 1990-1993. Reproduced from EC (1997).

pH 2.8. In the Southern Appalachian Mountains, cloud exposure at high elevation sites has been found to occur on 70% of the days and 30% of the time (Saxena and Lin, 1990). In addition 25% of all cloud events over a 4-year period had a pH \leq 3.1 with a minimum pH of 2.2 (DeFelice, 1997).

Recent air quality modeling indicates that eastern regional deposition of SO₄ and NO₃ is much larger than amounts measured in precipitation (CAPMON, NADP) would indicate (EC, 1997). This raises the real possibility that inputs of acidic deposition to some forests may be substantially higher than estimated.

2.2. OZONE

Composite national daily maximum 1 hour O₃ concentrations in the US decreased 15% between 1987 and 1996. The highest national composite 1 hour maximum was in 1988. Ozone levels have declined 10% since 1987 at 194 rural monitoring sites (EPA, 1998). In Canada, time series analysis identified a significant declining trend in daily maximum O₃ concentrations ranging from -0.05% to -0.08% yr⁻¹. Average days per year (1986-1993) exceeding the 82 ppb 1 hour National Ambient Air Quality Objective (NAAQO) ranged




Figure 4. Four-year mean wet deposition patterns of H⁺ (kg ha⁻¹yr⁻¹) in eastern North America for (a) 1980-1983 and (b) 1990-1993. Reproduced from EC (1997).

from 18 in southern Ontario, 3 in the Southern Atlantic Region (SAR) and 2 in the Lower Fraser Valley, British Columbia. (Olivotto, 1997). Mapping of percent change in SUM60 between 1985-1987 and 1994-1996 shows a significant decrease of 10-15% in much of the southeast US with decreases of 5-10% or 0-5% in the Lake States and southern Ontario/Quebec. Portions of the New England states and the SAR, however, had increases in SUM60 O₃ of between 0-10%. Smaller areas in the midwestern US showed increases of 10-15% SUM60 O₃ (Figure 5).

Ozone (photochemical smog) and dry-deposited N remain the most abundant air pollutants in U.S. Pacific Southwest forests (Fujioka *et al.*, 1998). In 1995, 99.3% of O₃ exceedances occurred during June to October. The number of adverse episodes in the Los Angeles-Long Beach and Riverside-San Bernardino end of the Southern California Air Basin (SoCAB) declined during 1986-1995. However, the differential between the western (urban) and eastern (urban/forested) ends increased (EPA, 1997). The most abundant reactive compound in the San Bernardino Mountains is still O₃, and ambient concentrations there are the highest in North America. Between November 2, 1991 and September 30, 1993, hourly concentrations exceeded the then-existing 120 ppb NAAQS on 235 days (22% of sampling days).




Figure 5. SUNM06 (May 1 - July 31; 08:00-19:59 hrs) ozone ppm-hrs calculated using Canadian (NAPS) and US (AIRS) monitoring data for (a) three-year average for 1994-1996 and (b) percent change between the periods 1985-1987 and 1994-1996. Source, T. Dann, Environment Canada-EPS.

3. Monitoring North American Forest Health

In the 1980's, the rationale and protocols for the North American monitoring programs changed due to studies that indicated that widespread changes in forest condition were occurring in North America and Europe (Rehfuess, 1981, Ulrich, 1984, Schutt and Cowling, 1985; McLaughlin, 1985; Fraser *et al.* 1985, Prinz, 1987). In Table I we have highlighted central features of both US, Canadian, and European forest monitoring networks, and some details of their structure and focus are discussed below.

3.1. THE UNITED STATES

The Detection Monitoring component of the US Forest Health Monitoring Program (FHM) (USDA, 1992) was implemented in 1996 on about 40% of approximately 4000 forested plots in the lower 48 states. A spatially and temporally systematic sample has been based on a national hexagonal grid (27 km between grid centres). The plot size and sampling frequency format (Table I) results in all circular 1 ha plots being sampled every 4 years with annual estimates of change being derived from smaller subplots (Smith and Mangold, 1996). This design yields: 1. a base-line condition the first year; 2. a direct measure of 1-year change for sub-plot samples after two years; and, 3. a direct measure of 4-year change after five years. Indicators (Table I) being measured in Detection Monitoring include lichen communities, ozone bioindicator plants, tree growth, crown condition, damage, mortality, regeneration, vegetation structure and plant diversity (Smith and Mangold, 1996).

3.2. CANADA

The Acid Rain National Early Warning System (ARNEWS) was initiated in 1984 (D'Eon and Power, 1989). ARNEWS consists of 150 rectangular, 10m X 40m plots with 2m x 2m subplots stratified by forest region. The system currently contains about 11,700 trees (dbh \geq 10cm) comprised of 3,500 hardwoods and 8,200 softwoods (Hall, 1995). Soil samples, foliar samples for nutrient analysis, and radial growth measurements are collected every five years adjacent to the plot (soil) or from numbered off-plot trees (foliage, increment cores). Variables are sampled on all plots at varying frequencies (Table I). Foliar symptoms, hardwood seed crop, pest conditions (type of foliar insect/disease, woody tissue insect/disease and severity) and tree condition (mortality, live crown height, crown damage, needle retention) are assessed seasonally and/or annually. Regeneration, saplings and ground vegetation (mosses, herbs, shrubs) are assessed biennially (D'Eon *et al.*, 1994).

3.3. JOINT UNITED STATES AND CANADIAN MONITORING

Out of growing concern over sugar maple decline in the early 1980's, the USDA Forest Service and the Canadian Forest Service joined in 1988 with state and provincial agencies to establish the North American Maple Project (NAMP). NAMP consists of a network of 233 sites extending across the natural range of sugar maple (*Acer saccharum* Marsh) (Millers *et al.*, 1991). NAMP sites are five-plot clusters of 20m x 20m plots on which all trees of 10 cm dbh or more are identified. Stands ranged from 50 to 150 years of age in 1988 and had variable levels of damage or crown dieback. Sites were paired where possible, one being tapped and the other untapped (no tapping or management during previous 5

TABLE I
Comparison of the United States Forest Health Monitoring (FHM), the Canadian Acid Rain National Early Warning System (ARNEWS), the North American Maple Project (NAMP) and the European (UN-ECE ICP) forest health monitoring programs.

	Monitoring Program			
	FHM	ARNEWS	NAMP	UN-ECE
No. plots	4000	150	166	20,345
Plot strat. ¹	grid 27km ²	forest region	na	grid 16km ²
Plot visits				
all	year 1/5	1	1	1
subset	1/3 year 2 1/3 year 4 1/3 year 5			
Associated studies	intensive site	FIGC	none	Level II
Tree				
diameter	E	5	E	X
height	E	5	X	X
dominance	E	5	X	X
form	X	5	X	X
crown height	X	5	X	X
crown closure	X	5	X	X
mortality	1-4	1	1	X
Crown				
condition	1-4	1	1	1
symptoms	1-4	1	X	1
insects/diseases	1-4	1	1	X
foliage	1-4	1	1	X
defoliation	1-4	1	1	1
woody	1-4	1	1	X
seed production	X	1	X	X
Ground				
regeneration	1-4	2	X	X
vegetation type	1-4	2	X	X
Pests	1-4	S	X	X

TABLE I

Continued

	Monitoring Program			
	FHM	ARNEWS	NAMP	UN-ECE
Nutrition				
foliar	X	5	X	X
organic soil	X	5	X	X
mineral soil	X	5	X	X
Metereology/Air Quality				
temperature	X	X	X	X
precipitation	X	X	X	X
SO ₄ /NO ₃	X	M	X	X
O ₃	B	P	X	X

FIGC : CFS Forest Indicators of Global Change

Sampling frequency: 5=5 yrs; 4=4yrs; 2=2yrs; 1=yearly; S=seasonally (several times per year); E=on establishment; X=not done

M: modelled; B: bioindicator plants; P: passive monitoring

1. Stratification of plots in network.

years). Variables assessed annually include defoliation, tree vigor, new damage to boles and crown condition (dieback, transparency).

The three forest health monitoring systems were designed to different protocols in order to address unique agency mandates. Each has specific strengths and weaknesses in responding to the monitoring of forest health response to changing air quality. In our view, none, by themselves, will enable forest managers or policy makers to relate effects to causes with the required degree of scientific certainty.

3.4. INSECT AND DISEASE PROBLEMS IN NORTH AMERICA

An important backdrop for consideration of pollution-induced stresses of North American forests is the extent of problems from insects and disease. A summary of the types of diseases, their approximate extent (where available) and the principal regions in which they have been detected, is given for the USA and Canada in Table II. Disease problems to forests are a major consideration in both Canada, where annual forest depletion due to disease during 1982-87 totaled 51 M cubic meters (NRCAN, 1996) and in the U.S.A., where the areal extent of insect problems on forests is 20.4 M hectares (USDA, 1997). With this background we consider the types of physiological effects induced by regional air pollution from two perspectives: (1) the relative magnitude and distribution of effects attributable to air pollution; and (2) the potential for stresses from air pollution to interact with biotic and abiotic stresses.

TABLE II
Estimates of forested areas affected by pests and pathogens in the U.S. and Canada in 1996.
Data are in millions (M) of hectares (where available) by region.

<u>United States¹</u>	
<u>Quantified</u>	
Dwarf mistletoe (11.7 M-W), fusiform rust (5.5 M-SE)	
Southern pine beetle (3.0 M-SE), spruce budworm (0.3 M-E+W)	
Mountain pine beetle (0.1 M-NW), gypsy moth (0.1 M-NE)	
<u>Unquantified</u>	
Beech bark disease (NE), butternut canker (NE)	
Hemlock woody adelgid (NE), dogwood (E)	
Anthraxnose (NE, SE), European pine shoot beetle (NE, NC)	
Total area affected > 20.4 M ha	
<u>Canada²</u>	
<u>Quantified</u>	
Aspen 2-leaf tier (3.9 M-W), eastern spruce budworm (1.2 M-E)	
Forest tent caterpillar (1.2 M-Canada), other insects combined (7.1 M-Canada, W)	
Other insects including area of beetle killed trees (4.6 M-Canada)	
¹ Source: USDA (1997)	
² Source: NRCAN (1997)	

4. Regional Air Pollution: North American Case Studies

Evidence of regional threats to forest health from air pollution in the U.S.A. led in the mid-1980s to the National Forest Response Program (Hertel and McKinney-McNeal, 1991), an integrated research program aimed at evaluating evidence and mechanisms of forest responses to acidic deposition and O₃. Initial integrated summaries were produced around 1990 in both the US (NAPAP, 1990) and Canada (Hall and Addison, 1991). The 1990 NAPAP synthesis included both studies of forest health (Barnard *et al.*, 1990) and evaluation of process level changes associated with regional air pollution (Shriner *et al.*, 1990). While very different in context (utilitarian vs ecosystem focus), the conclusions were similar in message - no clear cut evidence of a widespread forest decline, except for high elevation red spruce. Both analyses noted, respectively, uncertainty and evidence for pollution-induced changes in nutrient cycles and biotic and abiotic stress interactions that support longer term forest health. Canadian analyses in the NAPAP era, (Pearson and Percy, 1990) indicated that sugar maple decline was more severe and extensive than historically noted, and that acid deposition was a co-factor in a complex of contributing stressors. White birch decline along the east coast was circumstantially linked to acid fog. Critical loads were unavailable, but 46% of the Canadian land surface mapped was considered highly sensitive to acidification.

These first assessments have now been augmented by more recent syntheses of integrated studies (Eagar and Adams, 1992; Johnson and Lindberg, 1992; Fox and Mickler, 1995; Mickler and Fox, 1997) that were still in progress in 1990. Coupled with more recent results from Canadian studies (Hall, 1995; Hall *et al.*, 1997) and synthesis of the long-

standing ecosystem research on oxidant stress effects in the California ponderosa pine forests (Miller and McBride, 1998), additional valuable insights into the nature and extent of regional air pollution stress have been provided that we draw on here.

4.1. SOUTHEASTERN PINE FORESTS

The health of southern pine forests has been a subject of great interest and substantial controversy since significant growth declines were detected in the mid 1980's (Sheffield *et al.*, 1985 and Sheffield and Cost, 1987). The evidence of unexplained growth reductions came from analysis of FIA inventory data from unmanaged pine stands by the US Forest Service. With an area of approximately 25 M ha and annual stemwood sales of \$4.5 billion, the health of southern pines has been a major economic concern within the region. This concern translated into a large multiagency research effort aimed at quantifying causes of the apparent growth losses of the three predominant pine species in the region. The Southern Commercial Forest Research Cooperative (SCFRC), which was organized to address this task, conducted a series of regionally distributed controlled exposure studies with seedlings and saplings of loblolly pine (*Pinus taeda* L.), shortleaf pine (*P. echinata* Mill) and slash pine (*P. ellioti*) to evaluate the potential of O₃ and acidic deposition to affect southern pine growth. The summary of forest characteristics, biotic and abiotic stresses, and potential impacts of acidic deposition and ozone for southern pine forests combined information gathered in epidemiological and mechanistic studies over approximately 8 years of research (Fox and Micker, 1995).

A summary of the results of controlled exposure studies with seedlings and saplings of southern pines in open-topped chambers in the SCFRC indicated that significant growth reductions occurred with increased ozone exposure, but no consistent adverse effects of acidic deposition were detected (Teskey, 1995). Most studies reported small reductions in growth at ambient ozone, estimated at 2-5% per year, but these reductions were not typically detectable statistically within the experimental constraints of the experimental systems employed (Teskey, 1995). Contributing to this variability were genetics (multiple families with very different sensitivity levels), treatments imposed (interactions of ozone and acid deposition) and substantial year to year variability in ambient-based treatments imposed.

Additional analyses of the combined results of a wide range of controlled ozone studies with loblolly pine emphasized the role of high genetic variability in sensitivity in growth responses to ozone, particularly at near-ambient levels (Taylor, 1994). Using this much larger data set, Taylor (1994) estimated an ozone response threshold of 33 ppm hrs and a 6% growth loss per 100 ppm hrs of ozone exposure for sensitive families. The threshold for the population mean was similarly estimated at 97 ppm hrs with a 3% growth loss per 100 ppm hrs of cumulative exposure. At typical annual ambient ozone exposure levels of 50-200 ppm hrs (Taylor, 1994), estimated annual growth losses during a single season were 1-10% for sensitive families and 0-3% for the mean population.

Several studies have now indicated that the effects of ozone exposure are cumulative and approximately linear. The concept, which apparently extends to effects occurring over multiple seasons, is supported by studies of both loss of photosynthetic capacity (Richardson *et al.*, 1992) and foliage production and retention (Stow *et al.*, 1992, Kress *et al.*, 1992). Reduced canopy photosynthesis was noted at a cumulative exposure threshold of 100-150 ppm hrs and reduced foliar retention occurred at 150-230 ppm hrs. An important principal

established in these analyses was the carryover of inhibition of photosynthetic potential of ozone impaired foliage into subsequent growing seasons (Sasek *et al.*, 1991).

Studies of responses of more mature loblolly pine trees have included modeling-based estimates developed around changes in gross primary production of branches (Dougherty *et al.*, 1992); regional application of a physiologically-based model with regional ozone exposure data (Hogsett *et al.* 1993); and empirically derived measures of tree growth responses in the field (Zahner *et al.*, 1989, and McLaughlin and Downing, 1996). Results of these analyses for pines and selected studies with hardwoods are summarized in Table III and indicate an expected response range of 0-35% annual growth reduction for mature pines, depending on species X year and ozone level. Work by McLaughlin and Downing (1997) and Zahner *et al.* (1989) support the possibility that sensitivity of mature trees to drought may be enhanced by ozone at levels occurring in the field. Using an empirically derived model from weekly growth responses of mature loblolly pine trees to ozone in the field, McLaughlin and Downing (1996) indicated that ozone significantly increased the effects of soil moisture stress on stem growth. This concept is supported by studies with pine seedlings and saplings at ambient or near-ambient ozone levels which document increased transpiration (Lee *et al.*, 1990) or increased water stress (Flagler *et al.*, 1997). In addition several European studies with saplings and mature conifers (Skarby *et al.*, 1987; Wallin and Skarby, 1992; Maier-Maercker and Koch, 1992; Maier-Maercker, 1997), have reported increased transpiration and/or reduced stomatal control of water loss for conifers at ambient ozone levels. Collectively these studies suggest that stomatal closure measured at high ozone levels with seedlings in some controlled exposure studies, may not adequately represent the expected response to ambient exposures in the field, particularly where large trees are involved.

SCFRC studies indicated a low short-term risk to growth from ambient exposure to acidic deposition, and in fact growth was typically stimulated by near-ambient levels (Teskey, 1995). However, acidic deposition exerts its effects primarily through nutrient mobilization and depletion, and the development of such effects in controlled studies will depend on initial soil nutrient content and soil volume, factors which affect the capacity for nutrient depletion. Of greater relevance from the perspective of evaluating risk to southern pines, is the regional distribution of pine stands on poorly-buffered soils depleted of base cations by previous agricultural practices. Approximately 60% of southeastern pine soils are considered to be susceptible to accelerated cation leaching (Richter and Markewitz, 1995). Binkley *et al.*, (1989) have estimated that 10-15% of commercial southern pine forests may be currently limited by low base cation supply. Long term studies of nutrient cycles at a reference watershed in South Carolina indicate that approximately 80% of the exchangeable calcium (Ca) has been lost from the soil over the past three decades, with 50% of this loss due to uptake by vegetation and 50% due to leaching (Richter *et al.*, 1994). Base saturation declined from $\geq 55\%$ in 1962 to 20% in 1972, and to 10% in 1990. This decline was most apparent in shallow soil profiles (< 20 cm) where fine roots are concentrated (Richter and Markewitz, 1995). In considering the potential negative effects of Ca depletion from these soils, it should be noted that many of these same soils are N deficient and are receiving 4.5-14 kg N/ha/y, approximately 15-45% of annual stand requirements, from atmospheric sources (Richter and Markewitz, 1995). Accumulation of N in the upper soil horizons has the potential to stimulate growth at the risk of longer term cation depletion, as we will discuss later.

TABLE III
Some estimates of effects of ozone on growth of forest tree species at ambient ozone levels
based on studies with seedlings and mature trees.

SEEDLING STUDIES

<u>Species</u>	<u>Growth Reduction</u>	<u>Conditions</u>	<u>Reference</u>
Multiple	0-10%	Shoot growth	Chappelka and Samuelson, 1998
Southern pines	2-5%	Summary estimate - SCFRC field-chamber studies	Teskey, 1995
Loblolly pine		Synthesis - whole tree biomass	Taylor, 1994
	0-3% 1-10%	Mean response and sensitive family response to 50-200 ppm*h	
Hardwoods	13%	Response surface derived values	Reich, 1988
Conifers	3%	at 20 ppm*h	

MATURE TREE STUDIES

Loblolly pine	2-9%	Whole tree carbon model using branch chamber data (GA)	Dougherty <i>et al.</i> , 1992
Loblolly pine	0-13%	Average annual weekly responses to O ₃ and interactions of O ₃ and moisture stress, 5 years (TN)	McLaughlin and Downing, 1996
	0-5%	Annual O ₃ effect - no water stress	McLaughlin and Downing, 1996
	0-30%	Annual O ₃ effect - moderate water stress	McLaughlin and Downing, 1996
Loblolly pine	5-12%	Regional forest model simulation based on small tree exposures	Hogsett <i>et al.</i> , 1993
Shortleaf Pine	1% y ⁻¹	Empirically derived growth loss 38% cumulative not explained by climate and competition. Natural stands 20-45 yr (GA, SC, NC)	Zahner <i>et al.</i> , 1989
Shortleaf Pine	Climatic shift	Dendroclimatic analysis - Increase in non-climatic influences beginning in 1963. (GA) ¹	Grissino-Mayer and Butler, 1993
White Pine	O ₃ > climate	Dendroclimatic analysis show O ₃ influences growth more than climate (ME)	Bartholomay <i>et al.</i> , 1997
Hardwoods	2-17%	Regional simulation with canopy-stand model across moisture gradients	Ollinger <i>et al.</i> , 1997

¹Authors attributed shift to non-climatic physiological factors. We have detected a close relationship between the annual pattern of residual variance (not explained by climate) and annual variations in regional ozone exposure levels.

4.2. SOUTHWESTERN PINE FORESTS

In the San Bernardino Forest, ponderosa pine (*Pinus ponderosa* Dougl. Ex Laws), one the most important forest species in California (Olson, 1992), is exposed to the highest ozone concentration in North America by topographical, meteorological and climatic influences which combine to advect the Los Angeles urban pollution plume inland along the western mountain slopes (Miller, 1992). The resultant chronic exposure and ecosystem level changes in the San Bernardino Forest Ecosystem to over 50 years of high levels of O₃ and oxidant air pollution has produced perhaps the best example of hierarchical forest response to air pollutants (Miller *et al.*, 1982; Miller *et al.*, 1989; Miller, 1992; Miller *et al.*, 1996). Ozone injury to the mixed conifer forest has been occurring since at least the mid-1950's in the mountains of Southern California (Miller and Millecan, 1971) and since the 1970's in the Sierra Nevada (Pronos *et al.*, 1978). Miller *et al.* (1963) were first to identify the classic O₃ symptomology and chlorotic foliar mottle associated with the injury. Species most affected by photochemical oxidants (mainly O₃) have been ponderosa and Jeffrey (*Pinus jeffreyi* Grev. and Balf.) pines.

Injury to pine foliage resulted in a decline in the health of sensitive trees; accumulation of weakened trees then resulted in a heavy bark beetle kill. This necessitated extensive sanitation logging to remove declining pine trees. Unable to compete for nutrients, water and light, remaining sensitive individuals declined allowing for enhanced growth of more tolerant species (Miller *et al.*, 1982). Chronic effects of O₃ and N compounds continue today. The area has been used as a natural laboratory to study the air pollution effects on ecosystem functioning, including human values, and the 30 year San Bernardino Case Study is now summarized in Miller and McBride (1998).

During early survey years of 1968-1972 O₃ concentrations were extremely high (>80 ppb O₃ for 1300 h for a five-month year). Ponderosa and Jeffrey pine mortality ranged from 8-10% (2-2.5% per year). Ozone concentrations at high elevation sites remained above 50 ppb at night. An increase in extent of severe foliar injury indicated the beginning of the selective death of ponderosa pine (Miller, 1973). Weakened trees were attacked by bark beetles and elimination of ponderosa pine from the mixed conifer forest was predicted (Cobb and Stark, 1970). This survey led to the establishment of a large, multidisciplinary research team to investigate essential ecological processes underlying mixed conifer forest response to O₃. A hierarchical response, parallel to that observed with eastern white pine (*Pinus strobus* L.) (McLaughlin *et al.* 1982), was documented. Average (24 h) concentrations of 50-60 ppb O₃: induced foliar injury leading to premature abscission; needle fall resulted in reduced photosynthetic capacity; decreased nutrient availability to stressed trees; reduced production of carbohydrates; and diminishing tree vigor resulting in decreased height and radial growth (Miller *et al.*, 1982). At the ecosystem level, changes in decomposition patterns and altered nutrient cycling altered successional patterns in some species. This cycle of interactions triggered by O₃ on sensitive individuals significantly altered ecosystem processes and intra-community relationships. Unlike a catastrophic event such as fire, tree death and associated energy changes within the ecosystem occur gradually. Understanding of such changes in tree vigor, species composition, energy flow, nutrient cycling, and interactions between trophic levels is critical if cause-effect relationships are to be adequately evaluated.

One test of cause-effect linkage is mixed conifer forest response to gradually diminishing O₃ stress. Miller *et al.* (1989) reported an improvement (1974-1988) in foliar

injury index at all but the most exposed plots in the San Bernardino Mountains in response to declining O_3 in 1976-1991 (Davidson, 1993). Changes in visual injury were paralleled by improving tree growth patterns. Growth plots in a high- O_3 exposure area along the southern rim of the San Bernardino Mountains showed an approximate 40% growth slowdown during the period of highest ozone (1945-72) with a recovery of basal area growth to pre-1945 levels during the most recent period of improved air quality (Arbaugh *et al.*, 1998). The concurrence of long-term reduction in precipitation and high O_3 were concluded to be responsible for the growth pattern, and not O_3 alone. However, a significant lag in integration of improved canopy condition into improved basal area (BA) growth was identified. BA increases in O_3 -sensitive ponderosa pine during the same period (Miller *et al.* 1991) remained less than the BA increase for O_3 tolerant species. Only continued ecosystem monitoring will define the new baseline.

Evidence has existed for some time (Hoffer *et al.* 1981) that polluted air masses accumulating over central California are transported into the Sierra Nevada Mountains (Stohlgren and Parsons, 1987). Surveys recently completed in national parks in the region documented a progression of foliar symptoms on ponderosa pine related to aspect and elevation (Peterson *et al.* 1991). Standardization of an index to assess crown injury on ponderosa and Jeffrey pines, the Ozone Injury Index (OII), occurred in 1989. The Sierra Cooperative Ozone Impact Study (SCOIAS) and the Forest Ozone Response Study (FOREST) were among the first reports linking crown assessment, including both foliar injury and needle retention, with cumulative ambient O_3 indices. This represents a potentially reliable, cost effective approach for monitoring and forecasting O_3 injury in remote areas (Arbaugh *et al.* 1998).

Critical co-factors affecting tree response in SCOIAS/FOREST were site climate, history, tree size and age, N deposition and genetics. The unique role of N deposition in the mixed conifer forests of California has been well established (Bytnerowicz and Fenn, 1996). Unlike other case studies, most deposition occurs in summer in dry N form as acidic vapor, gaseous and particulate species. Deposition may be as high as 25-45 kg N ha⁻¹ yr⁻¹ and has resulted in localized N saturation in some mixed conifer stands (Bytnerowicz and Fenn 1996).

In summary, the effects of chronic exposure of these forests to acidic deposition and O_3 were detected on forest health through intensive process research built upon a historical record of individual tree response (Miller *et al.* 1996; Bytnerowicz *et al.* 1998). The multidisciplinary study succeeded in providing a comparative database on mixed conifer forest exposed to moderate and high levels of gas and particle deposition. Acidic pollutants, despite extreme sensitivity of leaf surface wax to acidic hydrolysis by HNO_3 (Bytnerowicz *et al.* 1998), were not implicated in a short-term response. The interaction of O_3 with periodic droughts (Miller *et al.* 1996), in the long-term, and summer N pulses (Bytnerowicz *et al.* 1998), in the very short-term, were judged to be the greatest threats to the health of southwestern pine forests.

4.3. EASTERN SPRUCE-FIR FORESTS

High levels of mortality of red spruce (*Picea rubens* Sarg.) in the Northern Appalachians (Weeks and Smoronok, 1988) and Fraser fir (*Abies fraseri* (Pursh) Pair.) in the Southern Appalachians (Dull *et al.*, 1988) during the 1970's and 1980s were accompanied by reductions in radial growth of red spruce at many high elevation sites across the region

(Johnson and Siccama, 1983; McLaughlin *et al.*, 1987). The timing and distribution of observed changes in growth suggested a possible linkage to acidic deposition and an integrated series of field and laboratory studies was conducted over the interval 1987-1991 by the Northern Spruce Fir Research Cooperative (NSFRC, Eagar and Adams, 1992). A synthesis of this work concluded that both field and laboratory data support " --that regional scale air pollution has played a significant role in the decline of red spruce in the Eastern United States" (Johnson *et al.*, 1992). This was based on regional and local analysis of the epidemiology and distribution of increased tree mortality and canopy deterioration (Peart *et al.*, 1992); the timing and distribution of radial growth decline of mature trees (Cook and Zedaker, 1992); and mechanistic studies in the field and laboratory which linked changes in carbon allocation (McLaughlin and Kohut, 1992) and winter injury (DeHayes, 1992) to levels of acidic deposition monitored within the region (Mohnen, 1992). These studies represent an excellent model of effective integration of regional monitoring of forest condition with mechanistic studies designed to better understand observed responses. In this case, multiple lines of investigation yielded new mechanistic insights linking acidic deposition as a predisposing agent to increased stress of red spruce forests by abiotic stress. These included both winter injury and nutrient depletion from foliage and soils. A summary of the principal trends in responses observed in field studies from the Northern Appalachians is depicted in Figure 6.

Increased winter injury to current year foliage was an important symptom of damage detected on red spruce in the Northeastern mountains in the late 1950s and early 1960's (Friedland *et al.*, 1984). As this damage was repeated frequently over the following years, clear linkages were found to reduced radial growth of both montane and coastal red spruce forests (Jagels, 1986 and Percy *et al.*, 1993) and increased mortality of montane trees (Johnson *et al.*, 1992). By the mid 1980's increased mortality levels had reduced red spruce as a component of the high elevation canopy by $\geq 50\%$ in the Adirondack and Green Mountains and by 25% further North in New Hampshire and Maine. A dendroclimatological analysis of the growth decline for trees from affected areas revealed that the occurrence of increased winter injury was a consequence of increased sensitivity of trees to winter temperatures, not unusually cold winters (Cook and Johnson, 1989). A series of misting experiments with acidic deposition under controlled laboratory conditions (see DeHayes, 1992) coupled with exclusion studies with branches (Vann *et al.*, 1992) and chambered saplings (DeHayes *et al.*, 1991) at high elevation field sites indicated that ambient levels of acidic mist could reduce the cold hardiness of red spruce by 3-10°C. This is an amount sufficient to substantially increase winter injury to this species at high elevation northeastern sites (Johnson *et al.*, 1992).

The growth decline of high elevation red spruce in the Southern Appalachians began typically around 1965, 5 years later than in the North, was confined to elevations above 1700 m, and was not accompanied by the very high red spruce mortality levels of the Northern Appalachians (McLaughlin *et al.*, 1987). However, canopy condition of red spruce declined significantly at both high and low elevation sites during 1985-89 (Peart *et al.*, 1992) and average red spruce mortality was consistently higher (ranging from 3-4% per year) at high elevation sites than intermediate and lower elevations (Nicholas, 1992). Standing dead red spruce comprised 8-17% of the stands at high elevation sites where attrition of dead stems was found to be remarkably high - 33% per year (Nicholas, 1992). While a notable loss in basal area of red spruce has not yet been reported at the southern Appalachian sites (Smith and Nicholas, 1998) mortality has occurred across all size classes




Figure 6.

Results of many different analyses revealed a shift in the health of high elevation red spruce around 1960 corresponding to increasing loading from acidic deposition in the Northern Appalachian Mountains (After Johnson *et al.*, 1992). In summary left panels (top to bottom) describe reduced radial growth at high and low elevations, increased terminal leader mortality, shifts in wood cation chemistry, and increased frequency of winter injury to foliage. The timing of these changes coincides with sharp regional increases in emissions of S and N (upper right) and shifts in growth climate relationships that were not attributable to unusual climate (lower right panels). See Johnson *et al.* (1992) for sources and details.

and available data indicate that recruitment levels are only approximately 50% of mortality levels (Nicholas, 1992).

Physiological studies in the field at both Southern and Northern Appalachian sites have shown that reduced cation availability associated with atmospheric deposition of strong anions reduces net carbohydrate production by foliage (McLaughlin *et al.* 1990, 1991, Amundson *et al.*, 1992; and Schaberg *et al.*, 1997). Reduced growth, decreased photosynthesis, increased respiration, and foliar cation depletion observed in the field were reproduced with red spruce seedlings under controlled conditions with ambient range (pH 3.0) acidic mist (McLaughlin *et al.*, 1993). Reduced Ca availability induced by both interference with root uptake of Ca from the soil and increased leaching of Ca from foliage by acidic mists appear to be an essential feature of observed responses in these studies (McLaughlin and Kohut, 1992). Changes in membrane-associated Ca have now been linked to changes in winter hardiness of red spruce (DeHayes *et al.*, 1997) and subsequent controlled studies have shown that acid mist exposures can alter membrane-associated Ca and reduce winter hardiness of red spruce by 10°C (DeHayes *et al.*, 1999).

Dendroecological studies in the South (McLaughlin *et al.*, 1987 and McLaughlin *et al.*, 1998) in parallel to those in the North (Cook and Johnson, 1989) indicate that reduced growth of mature trees is associated with sensitivity to warmer late summer temperatures. These responses suggest that carbohydrate depletion by increased dark respiration, a physiological symptom associated with Ca deficiency (Bangerth, 1979 and McLaughlin and Wimmer, 1999), may have contributed to observed growth losses. Winter damage to foliage has not occurred to any significant extent in the Southern Appalachians, although a less severe chlorotic stippling, which accumulates over winter with increasing needle age has been detected (Anderson *et al.*, 1991). The absence of severe foliage loss may explain the lower mortality generally observed in southern red spruce stands. On the other hand foliar and soil Ca levels are generally lower (Joslin *et al.*, 1994) and soil solution aluminum levels are substantially higher at southern sites where soil solution Ca:Al levels ≤ 0.50 occur frequently at some high elevation sites (Johnson *et al.*, 1991).

While most of the research in the spruce fir ecosystem has been directed at red spruce, significant mortality has also occurred in co-dominant and closely-related balsam fir (*Abies balsamea* (L.) Mill) in the North (Weeks and Smoronok, 1988) and Fraser fir in the South (Dull *et al.*, 1988). Observed declines of these species are regarded by most to be primarily a natural phenomenon linked to cohort senescence (fir waves) and insect damage from the balsam wooly adelgid (*Adelges fraserii*). However the influence of Ca deficiency on a wide variety of plant processes including both senescence and disease resistance (Poovich, 1988) suggests that consideration of the role of Ca deficiency in predisposing trees to these stresses is warranted (McLaughlin and Wimmer, 1999). In the South, growth declines of mature fir began about 1965 (Adams *et al.*, 1985) at the same time that adelgid damage was first detected at multiple mountain sites (Eagar, 1984). Physiological studies of Fraser fir indicate that increased dark respiration linked to reduced Ca and increased elevation, parallels symptoms noted for red spruce (McLaughlin *et al.*, 1997). In the North, aggrading stands of balsam fir grew more slowly at high elevations compared to those of the same age at lower elevations after 1960 (Reiners and Lang, 1979), indicating that this species has experienced increasing stress at high elevation sites during the same time interval over which red spruce has been adversely affected by acidic deposition.

4.4. NORTHEASTERN HARDWOOD FORESTS

Abnormally high levels of mortality have been recorded for several species of hardwood trees in the northeast since the 1980's. Most concern has centered on sugar maple forests across large areas in Ontario, Quebec, Vermont and New Hampshire.

Extensive crown dieback and mortality of sugar maple was first detected during the late 1970's and early 1980's. By 1982, decline symptoms were evident across 2,000 km² of sugar maple forest in Quebec alone (Roy *et al.*, 1985). A sustained recovery of these maples has occurred since then with dieback and mortality now reduced to historical levels (Hall *et al.*, 1997). Hall (1995) reported sugar maple had an annual mortality of 0.9% during 1987-1993 with tree condition showing little yearly change. Similarly NAMP data collected during 1985-96 in the US showed the condition of sugar maple trees (n>4,000) in the New England and New York region was good during 1988-1996. Mortality averaged between 0.9% and 1.9% per year (Stoyenhoff *et al.*, 1998). Crown condition improved with transparency declining from over 18% in 1988 to less than 12% in 1996. Dieback declined from circa 7% in 1988 to circa 6% in 1996. No difference was determined for the health of sugar maple growing in sugarbush and non-sugarbush stands. In Canada condition of sugar maple crowns on NAMP (62 sites) plots generally improved from 1988 to 1993, particularly in Quebec which had the highest level of dieback in 1988 (Lachance *et al.*, 1995). Dieback ranged from 5.3% to 8.2% during 1988-1993 while transparency ranged from 9.1% to 21.6%. As in the US, crown condition and mortality did not differ between sugarbush and non-sugarbush stands.

While no relationship was initially found between average NAMP cluster dieback/transparency and average five-year mean wet SO₄ or NO₃ deposition (Lachance *et al.*, 1995), subsequent analyses based on regional differences in soil buffering provided additional insights. In initial analyses relationships were tested using deposition values from air monitoring stations not collocated with plot clusters and data were averaged regionally into three zones (low, medium, high). In subsequent analyses atmospheric deposition rates were superimposed on soil acidification potential based on criticals load calculation. Exceedances of critical loads were calculated (Arp *et al.*, 1996) for ARNEWS plots (Figure 7), and these analyses indicated that the majority of the plots located north of the limestone region in extreme SW Ontario experienced high exceedances of the calculated critical loads (Hall *et al.*, 1997). Arp *et al.* (1996) also reported using NAMP data for Ontario, that areas of critical load exceedance had consistently higher levels of branch dieback than areas of no exceedance. Hall *et al.* (1997) state that within the Canadian Shield, attainment of a 50% reduction in the greatest critical load exceedance would require a reduction of SO₄ deposition rates to 12 kg ha⁻¹ yr⁻¹ (See Figure 2).

Linkage of tree response to soil sensitivity to acidic deposition is closely tied to soil supply capacity for base cations. This is apparent in the northern Appalachians, where nutrient deficiencies K and Mg were initially associated causally with sugar maple decline (Bernier and Brazeau, 1988a,b; Bernier *et al.* 1989). This was supported by responses to fertilization with K, which improved foliar nutrient status and growth rates in declining maples (Ouimet and Fortin, 1992), and fertilization with Ca and Mg, which led to improved tree vigor (Hendershot, 1991). Studies in mature stands have confirmed that ambient levels of acid deposition have accelerated the loss of base cations from soils (Foster *et al.*, 1992) leading to reduced soil Ca, Mg and K concentrations in sugar maple stands. These effects have been greatest where deposition is highest (Morrison *et al.*, 1996). Similarly, liming

studies in sugar maple stands in Pennsylvania have demonstrated restoration of more vigorous growth and canopy health (Long *et al.*, 1997). In some sugar maple soil systems efficient in retaining and recycling added nutrients, benefits of cation additions made to soils to offset nutrient imbalances may last several decades (Hendershot and Courchesne, 1994).

Sugar maple is also sensitive to changes in concentration of soil NO_3 and increasing N deposition to some N-saturated forests is expected to cause nutrient imbalances, with resultant effects on growth (Yin *et al.*, 1994). In addition despite the scientific and public focus on the role of acidic deposition, other co-occurring stressors of sugar maple forests may be equally important. The role of O_3 , for instance, has not been adequately investigated. Most of the sugar maple range (NAMP plot distribution) is located in areas of high seasonal ozone exposure and increases in canopy transparency have followed years with higher ozone levels, particularly in 1988 when SUM60 across the sugar maple range increased from 6,600 ppb hrs O_3 to over 30,000 ppb hrs O_3 (Pearson and Percy, 1997). Transparency also increased during 1988-1993 (Lachance *et al.*, 1995) at those plots where O_3 exceedances were highest. Visible foliar injury was reported on three indicator species at 59%, 34% and 39% of FHM plots in the region during 1994-1996 respectively (Stoyenoff *et al.*, 1998).

In addition, winter damage from deep freezing of exposed soil and roots in 1980 in Quebec contributed to sugar maple decline and complicated symptom interpretation. Followup studies on sapflow changes following development of canopy dieback amplified by winter cold treatments of soil (Bertrand *et al.*, 1996 and Robitaille *et al.*, 1995) indicate that the degree of loss of root function may exceed estimates of damage to canopies based on visual symptomology. For example sapflow losses of 39% were found on trees with only 9-13% canopy losses (Robitaille *et al.*, 1995). Such canopy losses have previously been considered normal for sugar maple, but may signal more substantive disruption of belowground processes.

During the early 1980's, the condition of paper birch (*Betula papyrifera* Marsh.) and mountain paper birch (*Betula cordifolia* Regel) growing along the Bay of Fundy coast in Canada also deteriorated significantly. Initial observations of *Betula* spp leaf browning and premature abscission were recorded by the Forest Insect and Disease Survey (FIDS). Subsequent investigation excluded insects and diseases as causal agents. Atmospheric stressors were considered candidates, particularly acidic fog and O_3 (Magasi, 1985).

Fogs were collected and found to deposit substantial amounts of acidity ($0.1\text{--}0.5 \text{ keq ha}^{-1}$), which had not previously been detected and which exceeded amounts deposited to the same stands by rain. The fog exposure duration at a nearby site ranged up to 1400 hrs (27% of the time) during the growing season. Significant correlations were determined between H^+ ion ($P < .001$), NO_3 ($P < .001$) concentrations and leaf browning (Figure 8). Continued investigation of the two taxa confirmed the relationship between fog frequency and foliar browning. All trees were affected during 1987-1990 followed by an apparent recovery in tree condition concurrent with historical (1915-1990) lows in fog frequency. Recovery was attributed to the shedding of dead branches and twigs during winter storms and the lack of initiation of new dieback due to low fog frequencies. The lower taxon *B. cordifolia* was found to be more prone to browning, slower to recover from dieback and more prone to mortality (Cox *et al.*, 1996).

It has been known for some time that certain mid- to high-elevation species in the northeast like red spruce and balsam fir are adapted to cool, moist environments and are




Figure 8. Percentage of Fundy coast *B. cordifolia* (▲) and *B. papyrifera* (■) trees with dieback symptoms in relation to hours of fog (visibility < 1 km) (●) at the St. John airport from 1982 to 1993. Reproduced from Cox *et al.* (1996).

immersed in clouds/fog for up to 44% of the growing season (Lovett *et al.*, 1982; Schemenauer, 1986). Comparison of northeastern marine-corrected mountain cloud and coastal fog chemistries by Kimball *et al.* (1988) indicated coastal fogs were up to three times higher in ionic concentrations and were lower in pH. NO_3 concentrations were especially high given the higher O_3 concentrations measured along the New England coast.

The potential for direct damage to vegetation from acid fog and cloud water is high and Maine coastal fogs have among the highest acidities (pH 2.8) ever recorded (Cape, 1993). When dry deposition is factored into total atmospheric deposition to birch leaf surfaces, very strong, localized acidities are likely (Wisniewski, 1982), accentuated by meteorologically-driven leaf surface wetting/drying cycles.

5. Integrating Effects of Ambient Pollutant Levels on Forest Processes

5.1. EFFECTS ON TREE PHYSIOLOGICAL PROCESSES

Despite differences in species, levels and combination of principal and secondary stresses, and primary pathways of effects among the four case study forests, effects of air pollutants on the resources required for growth and defense are often similar. In Figure 9, changes in the primary resources that regulate forest function: carbon production and allocation, water supply, and plant nutrition are depicted. Important features and implications of alterations in these resources are discussed below.

5.1.1. Effects on Carbon Production and Allocation

Reduced photosynthetic production in response to chronic pollution stress was found in all four systems examined. This occurred both as a result of changes in photosynthetic rate per unit of foliage mass (a shift in biochemical production potential) as well as from changes in the amount of leaf area produced or retained under chronic pollution stress. The mechanisms of effect, and foliage age classes involved were quite different among the

different forests, however. While premature loss of older needle classes has been observed with ozone exposures in both western and southern pine forests, loss of upper canopy foliage of northern hardwoods is most closely associated with nutrient depletion of the outer canopy due to both foliar leaching by acidic deposition, and reduced uptake of Ca and magnesium from poorly buffered soils (Heisey, 1995; Ellsworth and Liu, 1994).

In red spruce significant loss of foliage can be caused by winter injury to current year needles, which are both the most productive and the most sensitive age class. Sensitivity is enhanced by exposure to mists at acidity levels that occur frequently at high elevation sites in the Eastern US, and has been linked to increased frequency of terminal shoot dieback (LeBlanc and Raynal, 1990, and see Figure 6). It is also associated with low Ca retention and is an apparent consequence of foliar leaching of membrane-associated Ca (DeHayes *et al.*, 1997). Losses of foliar Ca by individual rain events may amount to 30% of the foliar Ca content (Joslin *et al.*, 1988). Much slower loss rates have been estimated for sugar maple (10-133 day half-life) suggesting biological non-significance (Scherbatskoy, 1989). However, Ca signaling in leaf physiological processes occurs at levels three to four orders of magnitude lower than those found in cell walls, and even small leaching losses from the small but active metabolic pools in foliage may significantly alter cell physiological functions (McLaughlin and Wimmer, 1999). Both reduced foliar Ca retention and reduced




Figure 9.

A summary of the physiological changes induced by ambient air pollution in the four case studies indicates multiple pathways of effects on carbon, water, and nutrient resources required for forest health. The net effect of these process-level effects has been multiple indications of reduced carbohydrate production by affected trees. Such changes typically lead to a shift in relative allocation of energy toward growth at the expense of reduced resistance to biotic and abiotic stresses.

Ca supply rates from soil appear to contribute to reduced foliar Ca and increased physiological dysfunction of red spruce foliage (McLaughlin and Kohut, 1992).

Both chronic O₃ exposure of pines (Adams *et al.*, 1990, McLaughlin *et al.*, 1982) and exposure of red spruce to acidic deposition in the field (Andersen *et al.*, 1991), reduce basipetal translocation of photosynthetically-fixed carbon from source foliage. Increased retention of photosynthate has also been associated with increased respiratory costs associated with repair of oxidant-stressed foliage and contributes to the reduced availability of carbohydrates for new growth (McLaughlin *et al.*, 1982). Reduced foliar carbon production (reduced Pn rate and leaf area) and reduced allocation away from production centers leads to the reduced root growth frequently associated with ozone exposure (Cooley and Manning, 1987). By contrast the effects of acidic deposition on root growth and survival can occur both from reduced carbon production above ground (winter injury and increased respiration of foliage) as well as from toxicity associated with aluminum mobilization in the soil solution (Cronan and Grigal, 1995). More typically acidic deposition affects root growth by decreasing Ca:Al ratios to levels which interfere with Ca uptake and impair root physiological function (Joslin *et al.*, 1992, and Godbold and Kettner, 1991).

5.1.1. Nutritional Effects of Changing N and Ca Availability

Changing nutrient availability induced by atmospheric deposition can also alter plant allocation of carbon in ways that increase sensitivity to other stresses. Such changes include increased deposition and accumulation of N (Aber *et al.*, 1989), reduced deposition of atmospheric sources of Ca (Hedin *et al.*, 1994), and reduced availability of Ca in soils (Likens *et al.*, 1998). Increased absorption of atmospheric sources of N and metabolic utilization of N by forest canopies (Hanson and Lindberg, 1991, Fowler *et al.*, 1998) as well as increasing N availability from chronic accumulation in surface soils has the effect of shifting tree growth and allocation of resources aboveground at a cost of reducing root mass and reducing root:shoot ratios (Persson and Madji, 1995, Schulze, 1989). Such changes can place forests at increasing risks of reduced uptake of water as well as other mineral nutrients.

Ca, on the other hand, has a direct and indirect role in allocation of resources to defense and repair in plants as well as influencing respiratory metabolism. Availability of Ca is particularly important for developing plants because it is not translocated between plant organs and because it is required for a wide variety of metabolic functions associated with growth, structural integrity of cells and membranes, local and interorganismal signaling, and defense and repair processes associated with disease resistance (Hepler and Wayne, 1985). Ca supply is therefore linked to transpiration rate in trees and can be reduced by drought as well as factors which reduce root vitality. Accelerated leaching of Ca from foliage exposed to acidic mists (Joslin *et al.*, 1988) and reduced availability of Ca in soil solutions in poorly-buffered soils exposed to strong anion inputs (Reuss and Johnson, 1986; Robarge and Johnson, 1992) can combine to limit Ca uptake and increase physiological stress in acidic soils.

5.1.3. Altered water status

Air pollution stress can increase water stress in forest trees by two principal pathways: alteration of transpiration through control of water flux from leaves at the membrane (substomatal) or stomatal level; or effects on water uptake related to changes in fine root biomass or root function. Much of our thinking regarding ozone and water stress has been

based on seedling studies in which water stress has been shown to accelerate stomatal closure in response to high ozone exposures (Tingey and Hogsett, 1985). This assumption led Ollinger *et al.* (1997) to assume a protective role of water stress in modeling growth responses of mature forests to ozone stress. However, in controlled studies with 3-yr-old beech trees the stomatal closure which occurred when well watered trees were exposed to ozone was impeded under dry conditions (Pearson and Mansfield, 1993) suggesting a more complex regulation of water loss under these conditions.

Larger forest trees typically have much lower ratios of leaf area to root mass than seedlings and have a larger stem volume which can provide an additional hydrologic reservoir during stress episodes (Kozlowski *et al.*, 1991). While stomatal conductances of large trees of some western species can be much lower than for seedlings (Yoder *et al.* 1994, and Grulke and Miller, 1994), Samuelson and Kelly (1997) have reported both higher conductances and higher ozone uptake within the canopies of large trees than by nearby seedlings in Tennessee. The capacity for ozone to cause stomatal closure as a consequence of water stress may be reduced for larger trees at the moderate water stress levels which limit many physiological processes. Both cell wall growth and protein synthesis, for example, are reduced at stress levels well below those required to cause stomatal closure (Hsaio, 1973).

Studies in the field at ambient levels now provide direct and indirect evidence that moderate ozone levels can increase water stress and reduce growth in larger trees (see McLaughlin and Downing, 1997). Reduced stomatal control of water loss from conifers documented in European studies (Maier-Maerker, 1997) documents one mechanism for this response, however reduced root function provides a potentially additive pathway that has not been adequately evaluated to date. On the other hand in the more arid San Bernardino Forest, drought increased resistance of older needles to ozone induced injury. This suppression of O₃ injury to older whorls of ponderosa and Jeffrey pines during drought years, was followed by O₃ injury and needle abscission upon return of favorable soil moisture (Miller *et al.*, 1996).

5.2. EFFECTS ON BIOGEOCHEMICAL CYCLES

The primary effects of current and projected future air pollutant loading on biogeochemical cycles and associated forest health are associated with three processes that operate concurrently in at least some soils in all of the four case studies we have examined. These include depletion of base cations due to increased mobilization and leaching from forest soils (Federer *et al.*, 1989), alteration of nutrient availability in soil solutions through mobilization of acidic cations, AL, Fe, Mn, and H, (Robarge and Johnson, 1992 and Ulrich and Matzner, 1986), and finally the buildup of soil N to levels that saturate surface soils (Aber *et al.*, 1989).

5.2.1. Cation Losses

Nutrient budgets for diverse forest stands now document the increased leaching of base cations associated with inputs of the strong anions SO₄ and NO_x to forest soils (Johnson *et al.* 1985; Federer *et al.*, 1989; Johnson and Lindberg, 1992; Richter *et al.* 1994; Likens *et al.*, 1996; and Likens *et al.*, 1998). In general, acidic deposition has been found to approximately double leaching rates that occur naturally (Joslin *et al.*, 1992 and Johnson and Todd, 1987) Long term depletion of base cations from both foliage and soils in the

eastern US has led to accelerated depletion of exchangeable cations from mixed hardwood forested watersheds and predictions of Ca deficiency in coming decades (Federer *et al.*, 1989). Similarly combined effects of forest growth, an important acidifying influence that is a natural part of cation accumulation by aggrading forests (Ulrich and Matzner, 1986), natural leaching and acidic deposition depleted 80% of the base cations from surface soils of a southern pine site in South Carolina over 30 years (Richter *et al.*, 1994). Because of the importance of forest growth in accumulating Ca and other cations, whole tree harvesting, which removes nutrient rich branches as well as boles, can accelerate ecosystem depletion of cation pools (Federer *et al.*, 1989).

5.2.2. Acid Cation Interference

Two properties of cation exchange in forest soils are pivotal to the capacity of acidic deposition to reduce physiological availability of cations: First, aluminum is mobilized more rapidly than Ca from exchange sites in the soil solutions when strong anions are deposited on poorly buffered soils (Reuss and Johnson, 1986); second strong binding capacity allows the acidic cations $Al > Mn > Fe > H$ to replace Ca and Mg from both physical and biological binding sites. The first property leads to domination of Al in soil solutions in high N and S deposition areas with poorly buffered soils. As Ca:Al ratios fall below 1.0, a wide variety of studies indicate that the risk of adverse effects to forest physiological function increase (Cronan and Grigal, 1995). Analysis of fine root chemistry in both declining red spruce (Shortle and Smith, 1988) and declining sugar maple stands (Adams and Hutchinson, 1992) indicate that low Ca:Al levels in fine roots are associated with reduced growth of trees in poorly buffered soils.

The high replacement potential of acidic cations for Ca, can result in displacement of this essential cation from the cell walls and membranes of fine roots resulting in and loss of membrane integrity and reduced root absorptive capacity (Zhao *et al.*, 1987, Stienen and Bauch 1988 and Schroeder *et al.*, 1988). In addition, Ca binding in cell walls, an important feature influencing the structural integrity of wood, can be reduced by acid cation substitution in xylem walls (Momoshima and Bondietti, 1990) thereby potentially reducing wood structural integrity (McLaughlin and Wimmer, 1999).

5.2.3. Nitrogen saturation

The long-term inputs of atmospheric sources of N to forest soils have begun to produce measurable changes in forest N cycles in industrialized regions. The phenomenon of N saturation occurs when availability of ammonium and nitrate in soils exceeds the combined plant and microbial demands (Aber *et al.*, 1989; Schulze, 1989). This process can occur naturally where N-fixing species dominate (Cole, 1992), but in other forest types it results from the combined addition of $\leq 40 \text{ kg ha}^{-1} \text{ y}^{-1}$ of anthropogenic N added to the 20-75 kg/ha/y produced annually by decomposition processes of coniferous forests and 50-175kg/ha/y similarly produced by deciduous forests (Aber *et al.*, 1989). Based on European studies Schulze *et al.* (1989) have suggested critical loading thresholds for N of 3-14 kg ha⁻¹ y⁻¹ for poorly buffered soils and 3-48 kg ha⁻¹ y⁻¹ for calcareous soils (note current wet deposition loading patterns for North America in Figure 3).

The net effect of N saturation is accumulation of the acidic nitrate ion in the soil profile, reduced Ca:Al ratios in soil solutions, reduced root mass and depth, and increased N export to streams. Evidences of N saturation are now becoming apparent in many different forest types including high elevation Southern Appalachian spruce-fir (Nodvin *et al.*, 1995), mid-elevation mixed hardwoods in North Carolina (Swank and Vose, 1997),

West Virginia (Gilliam *et al.*, 1996), Northern Hardwoods in both the US (Aber *et al.*, 1989, and Likens *et al.* 1998) and Canada (Foster *et al.*, 1992) as well as in areas of regionally high N deposition in California (Bytnerowicz and Fenn 1996). Nitrogen saturation has implications for long term stability of forest nutrient cycles because it can lead to breakdown of the cation- rich O horizon, and increased leaching of cations from deeper soil horizons (Shortle and Bondietti, 1992). Evidence suggests that the low soil Ca resulting from this process, affects root function, including uptake of N, thereby potentially contributing further to N accumulation in the soil (McLaughlin and Wimmer, 1999). Under these conditions forests can be expected to become increasingly dependent on recent litter for required Ca levels and roots to colonize shallower soil layers in the forest floor (Persson and Madji, 1995 and Joslin and Wolf, 1992). Another consequence of increasing N can be reduced growth of fine roots (Dougherty *et al.*, 1997).

5.2.4. Combined Effects on Forest Nutrient Cycling

So how significant are these combined processes in limiting nutrient availability to regional forests? Both comparative nutrient budgets across diverse forest types (Johnson and Lindberg, 1992) as well as variations in foliar and root nutrient levels across environmental gradients (Friedland *et al.*, 1988; Robarge *et al.*, 1989; and Joslin *et al.*, 1994) indicate that there are significant differences in plant-available nutrients, particularly the base cations, Ca and Mg associated with gradients in acidic deposition to forests. The Integrated Forest Study (IFS, Johnson and Lindberg, 1992) has provided nutrient budgets for at least two representatives for three of the four forest types we have addressed in this review. In Figure 10, we compare for spruce-fir, southern pines, and northern hardwoods, the distribution of Ca pools in vegetation, the forest floor, and soil exchangeable pools (Figure 10a) with the fluxes of Ca, including Ca uptake, leaching, and net increment (Figure 10b). These comparisons emphasize the large differences in nutrient capital both between and within systems, and they also demonstrate why the risks of acidic deposition to nutrient cycles can differ markedly within regions and between forest types. The lower exchangeable Ca pools at high elevation spruce fir forests (in both the Southern (GSM) and Northern (WF) Appalachian Mountains are readily apparent from these contrasts.

Long-term biomass modeling based on critical load exceedance calculations has predicted that forest productivity will decline in some regions with continued SO_4 and NO_3 deposition (Arp and Oja, 1992). Long-term (30 yrs) ecosystem research at Hubbard Brook, NH reported by Likens *et al.* (1996) has documented substantial losses of soil base cations due to acidic deposition. Important consequences for forest sustainability are predicted. Long-term research at the watershed level has also been instrumental in documenting SO_4 flux through nutrient-poor sugar maple forests indicating that soil buffering capacity of these forests has been depleted. On the other hand, there is also evidence at the watershed scale that reducing S emissions in the Northeastern US is beginning to have measurable effects of nutrient cycles of some forests. For example, at Turkey Lakes Watershed in central Ontario, monitoring of precipitation, nutrient cycling and streamflow chemistries from five basins has detected declining acidic concentrations in precipitation during 1981-1990 and this has resulted in lower concentrations in soil runoff and stream water. Decreases of up to 31% have reduced the flux of SO_4 in the soil (Foster and Hazlett, 1991), and will reduce rates of cation loss from these systems.

Foliar and fine root nutrient analyses provide the most important integrators of nutrient availability to forests. Analysis of foliar nutrient data suggest that foliar cation levels principally Ca and Mg are currently in the range of expected deficiency symptoms at wide




Figure 10.

Comparative Ca fluxes (a) and pools (b) for representative spruce, northeastern hardwood, and pine forests are important components of differences in sensitivity of these forests to acidic deposition (after Johnson, 1992). Forest types include: high elevation spruce-fir forests, Great Smoky Mountains (GSM) in TN, and Whiteface Mountain (WF) in New York; northern hardwood forests, Huntington Forest (HF) in NY and Turkey Lakes (TL) in Ontario; and southern loblolly pine forests in Durham, NC (Duke Forest (DF) and Oak Ridge, TN (OLP). Note the low net increment (a) and lower exchangeable soil pools (b) for high elevation spruce and northern hardwood forests compared to managed lowland pine forests

spread locations in the high elevation spruce fir forests (Friedland *et al.*, 1988, Joslin *et al.*, 1994). Reduced base cation levels have also been noted at higher elevations for the northeastern hardwood forest at Hubbard Brook (Likens *et al.*, 1998). Such data do not distinguish the relative roles of innately low soil fertility and any additive effects of acidic deposition on foliar nutrient uptake. However several lines of evidence suggest that acidic deposition had been an important contributing factor to reduced cation availability in areas of high atmospheric deposition: These include (1) accelerated leaching of base cations from foliage by acidic rain and mist events (Joslin *et al.*, 1988), (2) demonstrated physiological significance of relative small changes in the small, active pools of leaf Ca (McLaughlin *et al.*, 1993 and DeHayes *et al.*, 1997); (3) close linkages observed between input of SO_4 and NO_3 , elevated levels of Al in soil solutions and depression of Ca:Al ratios in soil solutions to levels (seasonal Ca:Al mean = 0.40, Johnson *et al.*, 1991) at which cation uptake by roots is inhibited; (4) shifting patterns of cation content of annual tree rings that indicate that substitution of acid cations for basic cations has occurred in recent decades in areas

receiving high deposition of atmospheric acids (Bondietti *et al.*, 1989, Bondietti *et al.*, 1990, and see Watmough, 1997); and finally, (5) Retrospective analysis of foliar cation levels at a Southern Appalachian spruce/fir site, which indicates that foliar levels of Mg and Ca have decreased by 50% and 33% respectively during the past twenty years (Shepard *et al.* 1995).

6. Implications of Changes in Forest Processes for Forest Health

From the utilitarian perspective the direct effects of air pollution on forest health that can be documented at present on North American forests may be considered to be relatively small compared to documentable effects of other types of stresses, such as fire and biotic diseases (see Table II). The most direct linkages can be made to the regional forest types we have chosen as case studies. With red spruce, sugar maple, white birch, and western pines, all predominantly older growth natural communities, the symptoms are reduced growth, weakened trees, increased susceptibility to stresses, and increased mortality. These are the classical indicators of the multicomponent decline complex in which air pollution can be a contributing factor (Manion, 1981). In two of the four systems, northeastern montane red spruce and southwestern pines in the San Bernardino forest, decreasing health had been translated into changes in stand structure and diminished basal area, meeting the utilitarian definition of health loss. With southern pines, a substantial component of which is managed commercial forest, data from controlled chamber studies, dendroecological studies, and models (see Table III) indicate that growth of mature pines is being reduced by current typical ambient ozone levels at annual rates that may vary from 0 to 10% per year. However, at present there are no indications of community-level changes attributable to ozone, and the biological effects of cation depletion of southern pine soils on health of these forests are still conjectural (NAPAP, 1998).

From the ecological perspective, however, we must ask how the process-level indicators of altered forest function from these few case studies relate to responses of these and other forest types to the range of biotic and abiotic stresses that normally regulate forest growth and development. In Figure 9 we summarized relationships among processes affected by ambient levels of air pollutants and changes in water, carbon, and nutrient resources that can be documented at those levels. These are of course the same media through which climate, competition, and disease affect forest processes and forest health. In the regional forest environment it is important to recognize that chronic stress from air pollution can only be expressed through altered sensitivity of forest physiological processes to natural biotic and abiotic stresses. Additionally, internal shifts in resource allocation strategies related to plant allometry (McLaughlin and Shriner, 1980), stage of growth (Mooney and Chu, 1974), nutrition (Clancy *et al.*, 1995) age (Waring, 1987), species type (Waring and Cobb, 1992) and stand level nutrient balance (Mattson and Addy, 1975) significantly influence plant allocation of resources among the competing demands of growth, defense, and reproduction and hence can alter forest sensitivity to insects and diseases.

Evidence from process level studies indicates that regional levels of air pollution stress measured in our case study forests are reducing carbon reserves, increasing water stress, and reducing nutrient availability. Carbon (energy) reserves are critical to plant defenses in many ways, including notably formation of defense chemicals that prevent damage and repair or replacement of damaged tissues after attack. The allocation of carbon to defense

in forest trees is a much lower priority than maintaining leaf and root function (Waring and Pittman, 1985), thus chemical and structural defenses may be more significantly reduced at the organ or whole tree level by stresses which reduce carbohydrate production levels. Root diseases of conifers, including littleleaf disease (*Phytophthora cinnamomia*) of shortleaf pine (Hepting, 1945) and *Armillaria* root rot (Wargo, 1972) which often develop as secondary pathogens following predisposing stresses (Manion, 1981) are enhanced by reduced carbohydrate supply to roots. The increased sensitivity of ponderosa pine to bark beetle attack in the San Bernardino Mountains following O₃ damage is a classical case of reduced defenses of forest trees being related to chronic air pollution stress (Cobb *et al.*, 1968).

Closely tied to reduced carbohydrate reserves, whatever the cause, is reduced root growth and function and ultimately increased water stress. Water stress frequently increases tree sensitivity to insect attack (Clancy *et al.*, 1995) although sensitivity can vary with level of stress (Lorio and Summers, 1986) and species type (Waring and Cobb, 1992). For example, susceptibility of conifers to insect herbivores was consistently increased by water stress (29 out of 31 studies) whereas broadleaves showed almost equal positive and negative responses (Waring and Cobb, 1992).

Significant strides in understanding the interplay of plant physiological responses to stress have come in the area of plant nutrition, specifically significantly expanded understanding of the role of Ca in regulating a wide range of physiological processes. These include intercellular and intraorganismic signaling (McAinsh and Hetherington, 1998 and Trewavas and Malho, 1997), chemical defenses (Pooviah, 1988 and Roberts and Harmon, 1992), and physical repair of cellular damage by disease (Dixon and Harrison, 1994). In addition to being essential for cell wall growth (Eklund and Eliasson, 1990), lignin is an important defense chemical against diseases such as damage to firs by the balsam wooly adelgid (Timmel *et al.*, 1986). Poor soil nutrition, including low Ca, P, K, and Mg has been recognized as a predisposing factor is sensitivity of Douglas fir to spruce budworm attack (Kemp and Moody, 1984). Stark (1965) lists 20 studies in which forest fertilization increases resistance of pines to insect attacks. Interestingly in 10 of 12 studies in which Ca was added alone or as a component of fertilization, reduced density or increased mortality of insect larvae was noted. Radiochemical studies suggest that Ca may be toxic to insects (see Stark, 1965). The loss and/or reduced availability of exchangeable soil Ca from soil reserves in response to long term deposition of S and N, thus has important implications for long term forest growth and forest ecosystem health.

Collectively the physiological effects of air pollutants would be anticipated to either predispose forest trees to other stresses, or amplify their negative effects. Only recently, have patterns of change over time (>10 yrs) at regional and national scales been statistically analyzed with the vigor required to attribute changes in forest health at the species level to individual and/or complexes of biotic and abiotic stressors, including air pollutants (Brandt *et al.*, 1999). For example, in North America the regional patterns of most frequent occurrence of major disease problems documented by forest surveys, are spatially consistent with the patterns of highest levels of ozone and acidic deposition as noted in Figure 11. Such a relationship is compatible with the role of strong anion deposition in depleting available pools of cations, notably Ca, from the relatively low soil pools in some Northeastern forests (see Figure 10). This supports the possibility that both carbohydrates and nutrient resources that influence forest health through altering tree resistance or recovery potential are relatively more affected by pollution stress in the same areas that insects and disease are apparently most problematic. Detailed statistical analyses might

reveal a statistical relationship between developmental trends and air quality that is beyond the scope of the more qualitative analyses presented here. While such patterns complicate analyses of cause and effect relationships on a regional basis, they also provide opportunities to examine component stresses from the perspective of process-level interactions involving carbon, water, and nutrient resources. Such analyses would be required to permit more robust tests of cause and effect relationships.




Figure 11.

The areas of the eastern US that currently have the greatest frequency of problems from forest insects and disease (USDA, 1997) also receive the highest deposition of S and N, and/or have the highest annual exposure to tropospheric ozone. Deposition data are area-integrated estimates at a state level from regional isopleths presented by Allen and Gholz (1995) based on summed wet deposition of S and N in kg/ha/y and relative ozone exposure for 1998-1999. The area-integrated ozone exposure index was > 850 for states in the very high category, 775-850 for the high category and 700-775 for the moderate category. By contrast, exposure indices for AL, ME, and IO were 500, 280, and 120, respectively.

7. Predicted Changes in Physical Climate and Effects on Forest Physiological Function

The linkages between changing global atmospheric chemistry and climate change have become solidified in recent years (Thomson, 1995, IPCC, 1998). On a global scale, temperatures are predicted to increase by 1-3.5°C by 2100. Spatial and temporal patterns

of precipitation are expected to change more variably, leading to increasing frequency of droughts in some regions. Stratospheric ozone depletion since 1980 is not predicted to recover until 2050, even under the most optimistic scenario (Madronich *et al.*, 1995). Resultant increases in biologically-effective ultraviolet radiation (UV-B, 280-315 nm) reaching forests may have important consequences for tree molecular biology (DNA), biochemistry, physiology, growth, regeneration and reproduction (Percy and Gordon, 1998)

as well as changes in plant succession and biodiversity (SCOPE, 1992). Ecological systems, including notably forests, are predicted to be significantly impacted by both the magnitude and variability of climatic shifts produced (IPCC, 1998). These responses, which include both positive and negative physiological effects that will influence process level changes induced by regional air pollutants, are reviewed very briefly below. Primary emphasis is on results from the Southeastern US, where an integrated series of studies has recently been completed (Mickler and Fox, 1997).

From a physiological perspective, the effects of increasing CO₂ and increasing temperature on net primary production, the effects of temperature on the frequency and severity of drought, and the effects of increasing N on nutrient uptake and allocation to shoots and leaching from soils appear to be the most compelling components of climate change that could significantly influence forest sensitivity to biotic and abiotic stresses. Predicted regional climate models for the Southeastern US associated with a doubling of CO₂ levels (Cooter, 1998), include a mean growing season temperature increase of 3.7-6.4°C with more variable and less certain effects on precipitation patterns.

A wide variety of controlled studies now indicate that increasing CO₂ will lead to significant increases in leaf photosynthetic rates and increased net primary production of forest trees (Wullschlegel *et al.*, 1997; Teskey *et al.*, 1998; and Norby *et al.*, 1999). Although very limited data are available for more mature forest trees, results to date suggest that positive responses are rather robust, in terms of absence of limitations by variations in supply of other resources (Wullschlegel *et al.*, 1997). Increasing temperature, on the other hand, affects many essential forest processes and in many different ways. Most notably the carbohydrate economy may be improved by extending the length of the growing season, but adversely impacted by the increased respiratory costs of temperature-driven metabolic processes. A 5°C temperature increase might increase dark respiration by 50% ($Q_{10} = 2$) and have a major impact on carbohydrate production and allocation. Based on branch studies, Teskey *et al.* (1998) found that a 2° C temperature increase, while it increased Pn at ambient CO₂, reduced the Pn response to increasing CO₂. In addition increasing temperature will increase evapotranspiration, thereby increasing potential limitations of water supply on the amount and allocation of growth (Schulze *et al.*, 1987).

Current global models used for predicting the distribution and condition of forests in response to global change (IPCC, 1998), consider increases in CO₂ and water availability, on forest growth, but, at present, do not consider potential implications of regional air pollutants on forest growth processes. Modeling results, for a 2X CO₂ scenario (Table IV) indicate that with climatic warming the potential forest cover in North America might increase by 24% (without) to 44% (with) CO₂ stimulation of growth. However, results from field chamber experiments provide evidence of the potential for O₃ and CO₂ co-exposure to offset predicted gains from elevated CO₂ and to enhance damage from O₃ (Kull *et al.*, 1996) and indicate a need for consideration of combined effects of these gases in future assessments.

Much of the simulated increases in Table IV occur as a result of increases in the distribution of temperate forests, which migrate northward, decreasing forest area in the conterminous USA and increasing potential forest cover in Canada and Alaska. By contrast, the modeling projects a significant increase in areas within each forest type in which declining growth would be expected, with 9 to 19% of the area being so affected if the positive effects of CO₂ are unconstrained. In the absence of positive CO₂ effects, climate change caused predicted productivity to decline over 45% to 89% of the total forested areas of North America.

TABLE IV

Predicted percentages of current forested area in North America that would lose or gain biomass with a doubling of pre-industrial atmospheric CO₂ levels. The effects of associated climate change without CO₂ potential physiological benefits of 2 x CO₂ are shown in the numerator of each response pair.¹

PERCENTAGE OF FORESTED AREA RESPONDING (-/+ CO₂)

	Biomass loss (-/+)	Biomass gain (-/+)
Boreal forest and Taiga tundra	25/5	49/83
Boreal Conifer	84/16	2/54
Temperate evergreen forest	55/8	12/66
Temperate mixed forest	44/4	26/95

¹Source: IPCC (1998), The Regional Impacts of Climate Change. These are considered conservative estimates of expected temperature effects due to delayed temperature equilibration to increasing CO₂ (IPCC, 1998). Total forest area predictions for North America are for a 42-44% gain with CO₂, and 21-24% gain in the absence of CO₂ effects.

The physiological basis of the IPCC (1998) model-predicted declines in forest health in the two times ambient CO₂ environment are based upon the effects of high temperatures on forest growth, and associated increases in drought, fire, and pathogens. In reality the spectrum of responses is likely to include both positive and negative responses in forest growth over time and across regions depending on the rate of change of the various climatic components and spatial variability in soils. In regional simulation of responses of loblolly pine stands to climate change in the Southeastern US (McNulty *et al.*, 1996), small increases in temperature (2°C) were found to have positive effects on Pn unless combined with reduced water supply. Non-linearity of expected responses is particularly likely in the case of N deposition (Aber *et al.*, 1989), which is envisioned to have initial stimulatory effects on forest growth on N-poor soils, such as for southern pines, followed by negative effects as cation depletion continues, leading potentially to N saturation.

Stimulation of growth regionally by N deposition, as has apparently occurred in some areas of Europe in recent decades (Sterba, 1996), also has potential parallel non-linear effects on global climate change. Modeling studies indicate that N-stimulation of growth in a CO₂-enriched environment could initially lead to significantly increased carbon and N storage in the soils, but at the risk of forests ultimately behaving as sources of NO₂ from microbial release of accumulated N (Norby, 1998 and Goulding *et al.*, 1998). The effects of warming on litter decomposition and release of accumulated nitrate represent an additional risk, particularly to N-saturated soils such as occur in the high elevation spruce-fir ecosystem (Joslin, and Johnson, 1997). Interestingly studies in Germany indicate that liming can drastically reduce N₂O release from forest soils fertilized by N (Brumme and Beese, 1992) a response that suggests important linkages between Ca depletion from soils and the ability of vegetation to utilize/retain added or excess soil N.

8. Synthesis and Recommendations

The case studies we have reviewed indicate that stress from regional scale air pollutants is affecting the physiological processes of forest tree species within diverse, widely distributed,

and regionally important forests. The processes affected and the stresses they produce are important to the way these forests grow and respond to biotic and abiotic stresses within their regional environments. If one measures forest health strictly in utilitarian terms, the level of effects to the near term economic health of the forests we have examined that can be clearly linked to specific effects of air pollutants may be considered small relative to the total suite of stress responses that affect forest health. However, if one views the potential importance of the processes affected in terms of implications for performance of these species within the framework of biotic and abiotic stresses that affect and are affected by the same resources, the potential ecological effects become potentially much larger, particularly over time. In this regard we consider the potential to be high for ambient ozone and acidic deposition stress to enhance the effects of drought, insects, disease, and high temperature in many situations, but to vary widely across the forest landscape just as these stresses vary. Similar conclusions have been reached in a synthesis of European physiological studies, which highlighted changes in carbohydrate production and allocation as critical to enhanced sensitivity of European forests to secondary stresses (Skarby *et al.*, 1998).

The well documented effects of ambient air pollutants on carbohydrate production and allocation, root growth, water uptake and control, and soil Ca and N levels indicate that regional atmospheric pollution could significantly enhance the adverse effects of global warming, particularly in the United States. While we expect increasing CO₂ to ameliorate air pollution effects on tree physiology by enhancing carbohydrate production and allocation to roots, biochemical interactions at the leaf level are less clear (Kull *et al.*, 1996). We expect the effects of increased temperature and increasing drought coupled with the concurrent effects of ozone, and acidic deposition (including N) on root vigor and root depth to increasingly place limitations on CO₂ induced enhancement of growth. Under these conditions, the enhancement of stress from drought, disease, and fire associated with climatic warming become more likely and the potential for interactions among the physical and chemical climate would be increased. Also to be considered in this mix are future trends in air pollutants. Reductions in S deposition are already underway and signal a measurable reduction in regional S deposition (Likens *et al.*, 1996). Predictions of future ozone levels depend not only on levels of N emissions, but also on regional climate, which will affect regional air patterns and the extent to which forests serve as sources for NO_x and volatile organic carbon. Current predictions for future ozone levels vary widely from likely reduced levels based on expected trends in N emissions (EPA, 1997) to increases of 30% in the next 25 years (NAS, 1992).

8.1. FUTURE MONITORING NEEDS

As Innes (1998) points out, despite the fact that most countries do not attribute damage to air pollutants, considerable care must be taken as absence of record may not mean that the particular cause of injury (e.g. air pollution) was not present. Development of better diagnostic techniques applicable to regional/national/multinational inventories of crown transparency and discoloration remain to be developed (Ferretti, 1998; Innes, 1998). The very nature of effects of chronic air pollution on forest ecological processes assures that attribution of causal relationships between air pollution and forest health will be very difficult without at least annual scale data on changes in system function. Because the carbon, water, and nutrient resources affected by air pollution are affected by many other stresses, it has required multidisciplinary team approaches to couple focused process level

studies aimed at documenting mechanisms of response with the patterns of change detected by survey approaches. The case studies we have presented indicate the value of designing surveys that consider both ecological and pollution gradients and make comparisons across ecologically analogous sites that tie in to hypothesized mechanisms of action. Such a gradient approach was developed by Legge *et al.* (1981) around a point source. It is now implemented in the "CFS Forest Indicators of Global Change Project" using sugar maple, white pine and red spruce ARNEWS/NAMP plots, with on-site process research, across gradients (2000 km) in acid deposition exceedance, O₃ exceedance and climate (temp. precip.) in eastern Canada.

Because of the temporal covariances among stressors in the environment, the development of a time series of multiple component environmental data provides an analytical framework within which to identify and model dominant independent variable and their interactions in affecting observed responses. As an example, the development of dendroecology as a method to evaluate annual variability in forest growth responses to physical and chemical climate has provided important tools to identify shifts in climate-growth relationships indicative of the timing and magnitude of increased external stresses, such as regional air pollution (McLaughlin *et al.*, 1987, Zahner *et al.*, 1989; Cook and Johnson, 1989, Petersen *et al.*, 1995). The development of additional analytical approaches such as multispectral scanning, which can be deployed with remote sensing technology, offers important new tools to obtain repeated measures of changes in canopy condition (Royle and Lathrop, 1997), canopy chemical changes such as lignin content indicative of soil nitrification rates (Wessman *et al.*, 1989), as well as providing indicators of more subtle shifts in leaf physiology, such as changes in photosynthesis and stomatal conductance (Carter, 1994) in response to environmental change. The statistical power provided by repeated measures of forest responses at interannual to annual scales can help provide important insights into the roles of both primary and secondary variables in controlling forest growth responses in a changing chemical and physical climate. This is substantiated by the retrospective statistical analysis of ARNEWS tree mortality (1985-1996) in relation to acidic deposition and other abiotic and biotic factors (Brandt *et al.*, 1999).

While we have focused primarily on physiological changes in this review, the development of quantitative indicators of tree condition also has great value in delineation of areas with outward evidences of changing forest health and mapping changes in those areas over time. Merging of monitoring and research expertise at the working levels will give impetus to the development, and implementation of more sensitive, cause-attributable diagnostic indicators. Successful examples of this concept in practice can now be found in both Europe (Swiss Federal Institute for Forest, Snow, and Landscape Research, Birmensdorf), and North America (Canadian Forest Service, Forest Health Network). In the past, monitoring and process-level research programs have tended to operate in parallel. However, monitoring data must be fully integrated with process studies if we are to more fully understand the influence of air pollutants and climate change on our forest health and long-term sustainability. The San Bernardino case study (Miller and McBride, 1998) remains the model of such a systematic approach to forest health.

In conclusion, our analyses indicate that changes in depth and vigor of root systems, shifts in pool sizes and allocation patterns of carbon, and changes in supply rates of N and Ca represent important shifts in ecological function that are currently occurring in diverse forest types across North America. The influence of these process level changes on future health of North American forests could be substantially increased in coming decades if projected changes in climate become a reality. Under such conditions, evaluating the

diverse influences of multiple stresses, operating in multiple combinations, will require monitoring programs that combine economic and ecological indicators of forest health (Olsen and Schreuder, 1997). Future monitoring programs should address multiple interactive hypotheses for causality with analytical tools that can adequately evaluate consistency, responsiveness, and mechanistic linkages of cause effect relationships across gradients in resource supply, both natural and manipulated. Such efforts should help forest ecologists to manage resources of North American forests in the future in ways that enhance both ecological and economic health.

References

- Aber, J. D., Nadelhoffer, K. J., Steudler, P., Melillo, J. M.: 1989, *BioScience* **39**, 378.
- Adams, C. M. and Hutchinson, T. C.: 1992, *Can. J. For. Res.* **22**, 1489.
- Adams, H. S., Stevenson, S. L., Blasing, T. J., DuVick, D. N.: 1985, *Env. Exp. Bot.* **25**, 315.
- Adams, M. B., Edwards, N. T., Taylor, G. E., Jr., Skaggs, B. L.: 1990, *Can. J. For. Res.* **20**, 152.
- Allen, E.R. and Gholz, H.L.: 1995, in S. Fox and R.A. Mickler (eds), *Impacts of Air Pollutants on Southern Pine Forests*, Springer, New York, p 83.
- Amundson, R. G., Hadley, J. L., Fincher, J. F., Fellows, S., Alscher, R. G.: 1992, *Can. J. For. Res.* **22**, 1605.
- Andersen, C. P., McLaughlin, S. B., Roy, W. K.: 1991, *Can. J. For. Res.* **21**, 455-461.
- Arbaugh, M. J., Miller, P. R., Carroll, J. J.: 1998, *Envir. Pollut.* **101**, 291.
- Arp, P.A. and Oja, T.: 1992, CEA No. 841-726, Joint CEA-CPPA-FC-EC Project Tech. Rep. Part Xvl.
- Arp, P.A., Oja, T., March, M.: 1996, *Can. J. For. Res.* **26**, 696.
- Bangerth, F.: 1979, *Ann. Rev. Phytopathol.* **17**, 97.
- Barnard, J., Lucier, A. A., Johnson, A. H., Brooks, R. T., Karnosky, D. F., Dunn, P. H.: 1990, NAPAP State of Science and Technology, National Precipitation Assessment Program.
- Bartholomay, G. A., Eckert, R. T., Smith, K. T.: 1997, *Can. J. For. Res.* **27**, 361.
- Beattie, B. L. and Whelpdale, D. M.: 1989, *Water Air Soil Pollut.* **46**, 45.
- Bernier, B. and Brazeau, M.: 1988a, *Can. J. For. Res.*, **18**, 754.
- Bernier, B., and Brazeau, M.: 1988b, *Can. J. For. Res.* **18**, 762.
- Bernier, M., Pare, D., Brazeau, M.: 1989, *Water, Air Soil Pollut.* **48**, 239.
- Bertrand, A., Robitaille, G., Boutin, R., Nadeau, P.: 1996, *Tree Physiol.* **15**, 775.
- Bertrand, A., Robitaille, G., Nadeau, P., Boutin, R.: 1994, *Tree Physiol.* **14**, 413.
- Binkley, D., Driscoll, C. T., Allen, H. L., Schoenenberger, P. and McAvoy, D.: 1989, *Acid deposition and forest soils: context and case studies in the Southeastern United States*, Springer-Verlag Ecological Studies **77**.
- Bondietti, E. A., Baes, C. F. III, McLaughlin, S. B.: 1989, *Can. J. For. Res.* **19**, 586.
- Bondietti, E. A., Momoshima, N., Shortle, W. C., Smith, K. T.: 1990, *Can. J. For. Res.* **20**, 1850.
- Brandt, J. P., Tamm, B. D., Hurley, J. E.: 1999, *Water Air Soil Pollut.* (submitted).
- Brook, J. R.: 1995, *Atmos. Environ.* **29**, 1795.
- Brumme, R. and Beese, F.: 1992, *J. Geophysical Res.* **97**, 12,851.
- Bytnerowicz, A., and Fenn, M.: 1996, *Environ. Pollut.* **92**, 127.
- Bytnerowicz, A., Percy, K., Riechers, Padgett, P., Krywult, M.: 1998, *Chemos.* **36**: 697.

- Cape, J.N.: 1993, *Environ. Pollut.* **82**, 167.
- Carter, G. A.: 1994, *Int. J. Remote Sensing* **15**, 697.
- Chappelka, A. H. and Samuelson, L. J.: 1998, *New Phytol.* **139**, 49.
- Clancy, K. M., Wagner, M. R., Reich, P. B.: 1995, pp. 118-180 in W. K. Smith and T. M. Hinckley (eds.), *Ecophysiology of Coniferous Forests*, Academic Press.
- Clarkson, D. T. and Hanson, J. B.: 1980, *Ann. Rev. Plant Physiol.* **31**, 239.
- Cobb, F. W., and Stark, R. W.: 1970, *J. For.* **68**, 147.
- Cobb, F. W., Wood, D. L., Stark, R. W., Parameter, J. R.: 1968, *Hilgardia* **39**, 141.
- Cole, D. W.: 1992, *Atmospheric Deposition and Forest Nutrient Cycling*, Springer-Verlag, pp. 150-151.
- Cook, E. R. and Johnson, A. H.: 1989, *Water, Air, Soil Pollut.* **48**, 127.
- Cook, E. R. and Zedaker, S. M.: 1992, pp. 192-231 in C. Eagar and M. B. Adams, *Ecology and Decline of Red Spruce in the Eastern United States*, Springer-Verlag.
- Cooley, D. R. and Manning, W. J.: 1987, *Environ. Pollut.* **47**, 95.
- Cooter, E. J.: 1998, pp. 15-54 in R. A. Mickler and S. Fox, *The Productivity and Sustainability of Southern Forest Ecosystems in a Changing Environment*, Springer-Verlag.
- Cox, R.M., Lemieux, G., Lodin, M.: 1996, *Can. J. For. Res.*, **26**, 682.
- Cronan, C. S. and Grigal, D. F.: 1995, *J. Environ. Qual.* **24**, 209.
- Davidson, A.: 1993, *Air Waste* **43**, 226.
- DeFelice, T. T.: 1997, *Atm. Res.* **43**, 325.
- DeHayes, D. H., Thornton, F. C., Waite, C. E., Ingle, M. A.: 1991, *Can. J. For. Res.* **21**, 1292.
- DeHayes, D. H.: 1992, pp. 295-337 in C. Eagar and M. B. Adams, *Ecology and decline of red spruce in the eastern United States*, Springer-Verlag.
- DeHayes, D. H., Schaberg, P. G., Hawley, G. J., Borer, C. H., Cumming, J. R., Strimbeck, G. R.: 1997, *Tree Physiol.* **17**, 687.
- DeHayes, D. H., Schaberg, P. G., Hawley, G. J., Strimbeck, G. R.: 1999, *Bioscience* (in review).
- D'Eon, S. P., and Power, M.: 1989, *The Acid Rain National Early Warning System (ARNEWS) Plot Network*, Forestry Canada Inf. Report PI-X-91, Petawawa.
- D'Eon, S. P., Magasi, L. P., Lachance, D., DesRochers, P.: 1994, *Canada's National Forest Health Monitoring Plot Network: Manual on Plot Establishment and Monitoring (Revised)*, Can. For. Serv. Inf. Rep. PI-X-117, Petawawa.
- Dixon, R. A. and Harrison, M. J.: 1994, *Annu. Rev. Phytopathol.* **32**, 479.
- Dougherty, P. M., Allen, H. L., Kress, L. W., Murthy, R., Maier, C. A., Albaugh, T. J., Sampson, D. A.: 1997, pp. 149-168 in R. A. Mickler and S. Fox, *The Productivity and Sustainability of Southern Forest Ecosystems in a Changing Environment*, Springer-Verlag.
- Dougherty, P. M., Teskey, R. O., Jarvis, P. G.: 1992, pp. 303-312 in R. B. Flagler (ed.), *Transactions: The Response of Southern Commercial Forests to Air Pollution*, Air and Waste Management Association.
- Dull, C. W., Ward, J. D., Brown, H. D., Ryan, G. W., Clerke, W. H., Uhler, R. J.: 1988, *Evaluation of spruce and fir mortality in the southern Appalachian mountains*, USDA Forest Service Southern Region R8-PR 13. 92 pp.
- Eagar, C.: 1984, *The Southern Appalachian Spruce-Fir Ecosystem: Its Biology and Threats*, National Park Service, Research/Resources Management Report SER-71.

- Eagar, C. and Adams, M. B.: 1992, *Ecology and Decline of Red Spruce in the Eastern United States*. Springer-Verlag. pp. 295.
- Eklund, L., and Eliasson, L.: 1990, *J. Exp. Bot.* **41**, 863.
- Ellsworth, D. S. and Liu, X.: 1994, *Can. J. For. Res.* **24**, 2118.
- Environment Canada (EC): 1997. Canadian Acid Rain Assessment Vol. 2: Atmospheric Science Assessment Report, Ottawa.
- EPA: 1996, *National Air Quality and Emissions Trends Report*, 1995. EPA 454/R-96-005.
- EPA: 1997, *National Air Pollutant Emission Trends, 1900-1906*, U.S. Environmental Protection Agency EPA-454/R-97-011.
- EPA: 1998, *National Air Quality and Emissions Trends Report*, 1996. EPA 454/R-97-013.
- Federer, C., Hornbeck, J. W., Tritton, L. M., Martin, C. W., Pierce, R. S., Smith, C. T.: 1989, *Environ. Manage.* **13**, 000.
- Ferretti, M.: 1998, *Chemos.* **36**, 1031.
- Flagler, R. B., Brissette J. C., Barnett, J. P.: 1997, pp. 73-92 in R. A. Mickler and S. Fox, *The Productivity and Sustainability of Southern Forest Ecosystems in a Changing Environment*. Springer-Verlag.
- Foster, N. W. and Hazlett, P. W.: 1991, pp. 179-191 In *Acid Deposition. Origins, Impacts and Abatement Strategies*, Springer-Verlag.
- Foster, N. W., Morrison, I. K., Yin, X. and Arp, P. A.: 1992, *Can. J. For. Res.* **22**, 1753.
- Fowler, D., Flechard, C., Skiba, U., Coyle, M. and Cape, J. N.: 1998, *New Phytol.* **139**, 11.
- Fox, S. and Mickler, R. A.: 1995, *Impact of Air Pollutants on Southern Pine Forests*. Springer-Verlag.
- Fraser, G. A., Phillips, W. E., Lamble, G. L., Hogan, G. D., Teskey, A. G.: 1985, *Can. For. Ser. Rep. E-X-36*. 43 p.
- Friedland, A. J., Gregory, R. A., Karenlampi, L., Johnson, A. H.: 1984, *Can. J. For. Res.*
- Friedland, A. J., Hawley, G. J., Gregory, R. A.: 1988, *Plant Soil* **105**, 189.
- Fujioka, F. M., Roads, J. O., Chen, S. C.: 1998, Climatology, in Miller, P. R. and McBride, J. (eds.), *Oxidant Air Pollution Impacts in the Montane Forests of Southern California: The San Bernardino Case Study*, Springer-Verlag, Heidelberg (in press).
- Gilliam, F. S., Adams, M. B., Yurish, B. M.: 1996, *Can. J. For. Res.* **26**, 196.
- Godbold, D. L. and Kettner, C.: 1991, *J. Plant Physiol.* **138**, 231.
- Goulding, K. W. T., Bailey, N. J., Bradbury, N. J., Hargreaves, P., Howe, M., Murphy, D. V., Poulton, P. R., Willison, T. W.: 1998, *New Phytol.* **139**, 49.
- Grissino-Mayer, H. D. and Butler, D. R.: 1993, *Southeast. Geogr.* **1**, 65.
- Grulke, N. E. and Miller, P. R.: 1994, *Tree Physiol.* **14**, 659.
- Hall, J. P.: 1995, *For. Chron.* **71**, 607.
- Hall, J. P. and Addison, P. A.: 1991, *Response to air pollution: ARNEWS assesses the health of Canada's forests*, For. Can. Sci. Dir. Inf. Rep. DPC-X-34, Ottawa, 42 pp.
- Hall, J. P., Bowers, W., Hirvonen, H., Hogan, G., Foster, N., Morrison, I., Percy, K., Cox, R., Arp, P.: 1997, 1997 Canadian Acid Rain Assessment: Vol. 4 The Effects on Canada's Forests, Natural Resources Canada, 47 pp.
- Hanson, P. J. and Lindberg, S. E.: 1991, *Atmos. Envir.* **25**, 1615.
- Hendershot, W. W.: 1991, *Fert. Res.* **27**, 63.
- Hendershot, W. W. and Courchesne, F.: 1994, *Can. J. For. Res.*, **24**, 609.
- Hedin, L. O., Granat, L., Likens, G. E., Bulshand, T. A., Galloway, J. N., Butler, T. J., Rodhe, H.: 1994, *Nature* **367**, 351.
- Heisey, R. M.: 1995, *Water Air Soil Poll.* **82**, 675.

- Hepler, P. K., and Wayne, R. O.: 1985, *Ann. Rev. Plant Physiol.* **36**, 397.
- Hepting, G. H.: 1945, *Phytopath.* **35**, 106.
- Hertel, G. D. and McKinney-McNeal: 1991, *USDA For. Serv. Agric. Inf. Bull.* **622**, 19.
- Hinckley, T., Ford, D., Segura, G., Sprugel, D.: 1992, Key processes from stand to tree level in Wall, G. (ed.), *Implications of climate change for Pacific Northwest Forest Management*. University of Waterloo, Department of Geography paper 15, Waterloo, pp. 33-43.
- Hoffer, T. E., Miller, D. J., Farber, R. J.: 1981, *Atmos. Environ.* **15**, 1935.
- Hogsett, W. E., Herstrom, A. A., Laurence, J. A., Lee, E. H., Weber, J. E., Tingey, D. T.: 1993, Proceedings of the 4th US/Dutch International Symposium: Comparative Risk Analysis and Priority Setting for Air Pollution Issues. Air and Waste Management Association.
- Hsaio, T. C.: 1973, *Ann. Rev. Plant Physiol.* **24**, 519.
- Innes, J. L.: 1993, *Forest Health: Its Assessment and Status*, CAB International, 677 pp.
- Innes, J. L.: 1998, *Chemos.* **36**, 1025.
- Intergovernmental Panel on Climate Change (IPCC): 1998, *The Regional Impacts of Climate Change. An Assessment of Vulnerability*. pp. 267-279 in R. T. Watson, M. C. Zinyowooa, R. H. Moss, and D. J. Dokken (eds.), Cambridge University Press.
- Jagels, R.: 1986, *I.A.W.A. Bull.*, 299.
- Jeffries, D. S. (ed.): 1997, *1997 Canadian Acid Rain Assessment: Vol. 3 Aquatic Effects*. Environment Canada, Burlington.
- Jenkins, A. F.: 1997, *J. For.*
- Johnson, A. H. and Siccama, T. G.: 1983, *Envir. Sci. Technol.* **17**, 294A.
- Johnson, A. H., McLaughlin, S. B., Adams, M. B., Cook, E. R., DeHayes, D. H., Eager, C., Fernandez, I. J., Johnson, D. W., Kohut, R. J., Mohnen, V. A., Nicholas, N. S., Peart, D. R., Schier, G. A., White, P. S.: 1992, pp. 385-411 in C. Eagar and M. B. Adams, *Ecology and Decline of Red Spruce in the Eastern United States*. Springer-Verlag.
- Johnson, D. W. and Todd, D. E.: 1987, *Plant Soil* **102**, 99.
- Johnson, D. W. and Lindberg, S. E.: 1992, *Atmospheric deposition and forest nutrient cycling*. Springer-Verlag.
- Johnson, D. W., Richter, D. D., Lovett, G. M., Lindberg, S. E.: 1985, *Can. J. For. Res.* **15**, 773.
- Johnson, D. W., Van Miegroet, H., Lindberg, S. E., Todd, D. E., Harrison, R. B.: 1991, *Can. J. For. Res.* **21**, 769.
- Joslin, J. D., McDuffie, C. M., Brewer, P. F.: 1988, *Water, Air, and Soil Pollut.* **39**, 355.
- Joslin, J. D. and Wolfe, M. H.: 1992, *Can. J. For. Res.* **22**, 893.
- Joslin, J. D. and Wolfe, M. H.: 1994, *Soil Sci. Soc. Am. J.* **58**, 1572.
- Joslin, J. D., Kelly, J. M., Van Miegroet, H.: 1992, *J. Env. Qual.* **21**, 12.
- Joslin, J. D., and Johnson, J. W.: 1997, pp. 571-590 in R. A. Mickler and S. Fox, *The Productivity and Sustainability of Southern Forest Ecosystems in a Changing Environment*. Springer-Verlag.
- Joslin, J. D., Wolfe, M. H., Garten, C. T.: 1998, pp. 557-570 in R. A. Mickler and S. Fox (eds.), *The Productivity and Sustainability of Southern Forest Ecosystems in a Changing Environment*. Springer-Verlag.
- Kemp, W. P. and Moody, U. L.: 1984, *Environ. Entomol.* **13**, 1291.
- Kimball, K. D., Jagels, R., Gordon, G. A., Weathers, K. C., Carlsile, J.: 1988, *Water Air Soil Pollut.* **39**, 383.

- Kolb, W. E. et al.: 1994, *J. For.* **92**, 10.
- Kozlowski, T. T., Kramer, P. J., Pallardy, S. G.: 1991, *The Physiological Ecology of Woody Plants*, Academic Press.
- Kress, L. W., Allen, H. L., Mudano, J. E., Stow, T. K.: 1992, *Environ. Toxicol. Chem.* **11**, 1115.
- Kull, O., Sober, A., Coleman, M. D., Dickson, R. E., Isebrands, J. G., Gagnon, Z., Karnosky, D. F.: 1996, *Can. J. For. Res.* **26**, 639.
- Lachance, D., Hopkin, A., Pendrel, B., Hall, J. P.: 1995, Health of Sugar Maple in Canada: results from the North American Maple Project, 1988-1993, Canadian Forest Service Report ST-X-10, 27 pp.
- LeBlanc, D. C. and Raynal, D. J.: 1990, *Can. J. For. Res.* **20**, 1415.
- Lee, W. S., Chevone, B. I., Seiler, J. R.: 1990, *Water Air Soil Pollut.* **51**, 105.
- Legge, A. H., Jaques, D. R., Harvey, G. W., Krouse, H. R., Brown, H. M., Rhodes, E. C., Nosal, M., Schelhase, H. U., Amundson, R. G., Walker, R. B.: 1981, *Water Air Soil Pollut.* **15**, 77.
- Likens, G. E., Driscoll, C. T., Buso, D. C.: 1996, *Science* **272**, 244.
- Likens, G. E., Driscoll, C. T., Buso, D. C., Siccama, T. G., Johnson, C. E., Lovett, G. M., Fahey, T. J., Reiners, W. A., Ryan, D. F., Martin, C. W., Bailey, S. W.: 1998, *Biogeochem.* **41**, 89.
- Long, R. P., Horsley, S. B., and Lija.: 1997, *Can. J. For. Res.* **27**, 1560.
- Lorio, P. L., Jr. and Sommers, R. A.: 1986, *Tree Physiol.* **2**, 301.
- Lovett, G. M., Reiners, W. A., Olson, R. K.: 1982, *Science* **218**, 1303.
- Madronich, S., McKenzie, R. L., Caldwell, M., Bjorn, L. O.: 1995, *Ambio* **24**, 143.
- Magasi, L. P.: 1985, *Forest pest conditions in the Maritimes in 1985*, Can. For. Sev. Inf. Rep. M-X-159, Fredericton.
- Maier-Maercker, U.: 1997, *Trees* **11**, 229.
- Maier-Maercker, U. and Koch W.: 1992, *Trees* **7**, 12.
- Manion, P. D.: 1981, *Tree Disease Concepts*, Englewood Cliffs, NY, Prentice Hall.
- Mattson, W. J. and Addy, N. D.: 1975, *Science* **190**, 515.
- McAinsh, M. R. and Hetherington, A. M.: 1998, *Perspectives* **3**, 22.
- McLaughlin, S. B.: 1985, *J. Air Poll. Control Assoc.* **35**, 516.
- McLaughlin, S. B., McConathy, R. K., Duvick, D., Mann, L. K.: 1982, *For. Sci.* **28**, 60.
- McLaughlin, S. B. and Shriner, D. S.: 1980, *Plant Disease Vol. 5*. Academic Press. pp. 407-431.
- McLaughlin, S. B. and Downing, D. J.: 1996, *Can. J. For. Res.* **26**, 670.
- McLaughlin, S. B. and Downing, D. J.: 1997, pp. 207-230 in R. A. Mickler and S. Fox (eds.), *The Productivity and Sustainability of Southern Forest Ecosystems in a Changing Environment*. Springer-Verlag.
- McLaughlin, S. B., Downing, D. J., Blasing, T. J., Cook, E. R., Adams, H. S.: 1987, *Oecologia* **72**, 487.
- McLaughlin, S. B., Andersen, C. P., Edwards, N. T., Roy, W. K., Layton, P. A.: 1990, *Can. J. For. Res.* **20**, 485.
- McLaughlin, S. B., Andersen, C. P., Hanson, P. J., Tjoelker, M. G., Roy, W. K.: 1991, *Can. J. For. Res.* **21**, 1234.
- McLaughlin, S. B., Tjoelker, M. G., Roy, W. K.: 1993, *Can. J. For. Res.* **23**, 380.
- McLaughlin, S. B. and Kohut, R.: 1992, pp. 338-384 in C. Eagar and M. B. Adams (eds.), *Ecology and Decline of Red Spruce in the Eastern United States*. Springer-Verlag.

- McLaughlin, S. B., Joslin, J. D., Robarge, W., Stone, A., Wimmer, R., Wullschlegel, S. D.: 1998, pp. 255-277 in R. A. Mickler and S. Fox (eds.), *The Productivity and Sustainability of Southern Forest Ecosystems in a Changing Environment*. Springer-Verlag.
- McLaughlin, S. B. and Wimmer, R.: 1999, *New Phytologist* (in press).
- McNulty, S. G., Vose, J. M., Swank, W. T.: 1996, *Ambio* **25**, 449.
- Mickler, R. and Fox, S.: 1997, *The Productivity and Sustainability of Southern Forest Ecosystems in a Changing Environment*. Springer-Verlag.
- Miller, P. R.: 1992, Mixed conifer forests of the San Bernardino Mountains in Olson, R. K., Binkley, D., Bohm, M. (eds.), *The Response of Western Forests to Air Pollution*, Springer-Verlag, New York. Pp. 461-497.
- Miller, P. R.: 1973, pp. 101-117 in J. A. Naegele, *Air Pollution Damage to Vegetation*. Advances in Chemistry Series 112. Am. Chem. Soc., Washington.
- Miller, P. R., Chow, J., Watson, J. G.: 1996, Assessment of Acidic Deposition and Ozone Effects on Conifer Forests in the San Bernardino Mountains, California Air Resources Board Report.
- Miller, I., Lachance, D., Burkmann, W. G., Allen, D.C.: 1991, North American maple decline project: organization and field methods, USDA Forest service Gen. Tech. Rep. NE-154.
- Miller, P. R. and McBride, J. (eds.): 1998, *Oxidant Air Pollution Impacts in the Montane Forests of Southern California: The San Bernardino Case Study*, Springer-Verlag, Heidelberg.
- Miller, P. R., McBride, J. R., Schilling, S. L., Gomez, A. P.: 1989, pp. 309-324 in R. K. Olson and A. S. Lefohn, *Effects of Air Pollution on Western Forests*. Air and Waste Manage. Assoc., Pittsburgh.
- Miller, P.R., and Millecan, A.A.: 1971, *Plant Disease Rep.* **55**, 555.
- Miller, P. R., Parmeter, J. R., Taylor, O. C., Cardiff, E. A.: 1963, *Phytopath* **53**, 1072.
- Miller, P. R., Chow, J., Watson, J. (eds.): 1996, *Assessment of Acidic Deposition and Ozone Effects on Conifer Forests in the San Bernardino Mountains*, California Air Resources Board Report NTIS 97115612, Sacramento.
- Miller, P. R., Taylor, O. C., Wilhour, R. G.: 1982, *Oxidant air pollution effects on a western coniferous forest ecosystem*. US EPA Report No. EPA-600/D-82-276.
- Millers, I., Lachance, D., Burkman, W. G., Allen, D. C.: 1991, *North American Sugar Maple Decline Project: Organization and Field Methods*. U.S. Dep. Agric. For. Serv., Radnor.
- Mohnen, V. A.: 1992, pp. 64-124 in C. Eagar and M. B. Adams (eds.), *Ecology and Decline of Red Spruce in the Eastern United States*. Springer-Verlag.
- Momoshima, N. and Bondietti, E. A.: 1990, *Can. J. For. Res.* **20**, 1840.
- Mooney, H. A. and Chu, C.: 1974, *Oecologia* **14**, 295.
- Morrison, I.K., Fournier, R.E., Hopkin, A.A.: 1996, *Proceedings 16th IUFRO International Meeting for Specialists in Air Pollutant Effects on Forest Ecosystems*, Fredericton, 402 pp.
- NAPAP: 1990, *National Acid Precipitation Assessment Program*, State of Science/Technology. Report Series.
- NAPAP: 1998, *National Acid Precipitation Assessment Program Biennial Report to Congress: An Integrated Assessment*, Silver Spring, MD.

- NAS: 1992, *Rethinking the Ozone Problem in Urban and Regional Air Pollution*. National Academy of Sciences. National Research Council. National Academy Press. 500 pp.
- Natural Resources Canada (NRCAN): 1996, *1995 Compendium of Canadian Forestry Statistics*. Canadian Council of Forest Ministers, Ottawa. 202 pp.
- Natural Resources Canada (NRCAN): 1997, *1996 Compendium of Canadian Forestry Statistics*. Canadian Council of Forest Ministers, Ottawa. 234 pp.
- Natural Resources Canada (NRCAN): 1998, *The State of Canada's Forests*, 109 pp.
- Nicholas, N. S.: 1992, *Stand Structure, Growth, and Mortality in Southern Appalachian Spruce-Fir*. Ph.D. Dissertation, Virginia Polytechnic Institute and State University, 176 pp.
- Nodvin, S. C., Van Miegroet, H., Lindberg, S. E., Nicholas, N. S., Johnson, D. W.: 1995, *Water, Air, and Soil Pollut.* **85**, 1647.
- Norby, R. J.: 1998, *New Phytol.* **139**, 189.
- Norby, R. J., Wullschlegel, S. D., Gunderson, C. A., Johnson, D. W., Ceulemans, R.: 1999, *Plant Soil Envir.* **22**, (in press).
- O'Laughlin, J., Livingston, R. L., Thier, R., Thornton, J., Toweill, D. E., Morelan, L.: 1994, *J. Sust. For.* **2**, 65.
- Olivotto, C. (ed.): 1997, *Canadian 1996 Nox/VOC Science Assessment: Summary for Policy Makers*, Environment Canada, Toronto.
- Ollinger, S. V., Aber, J. D., Reich, P. B.: 1997, *Ecol. Appl.* **7**, 1521.
- Olsen, A. R. and Schreuder, H. T.: 1997, *Envir. Ecol. Stat.* **4**, 167.
- Olson, R. K.: 1992, Physiography of forest types in Olson, R. K., Binkley, D. and Bohm, M. (eds.), *The Responses of Western Forests to Air Pollution*, Springer-Verlag, New York. pp. 7-40.
- Ouimet, R. and Fortin, J.-M.: 1992, *Can. J. For. Res.* **22**, 699.
- Peart, D. R., Nicholas, N. S., Zedaker, S. M., Miller-Weeks, M., Siccama, T. G.: 1992, pp. 125-191 in C. Eagar and M. B. Adams (eds.), *Ecology and Decline of Red Spruce in the Eastern United States*, Springer-Verlag.
- Pearson, M. and Mansfield, T. A.: 1993, *New Phytologist* **123**, 351.
- Pearson, R. G. and Percy, K. E.: 1990, *The 1990 Canadian Long-range Transport of Air Pollutants and Acid Deposition Assessment Report: part 5 Terrestrial Effects*, Federal/Provincial Research and Monitoring Coordinating Committee (RMCC), 105 pp.
- Pearson, R. G. and Percy, K. E.: 1997, Canadian 1996 NO_x/VOC Science Assessment: Report of the Vegetation Objective Working Group. Canadian Council of Ministers of Environment (CCME) Multistakeholder NO_x/VOC Science Plan, pp.
- Percy, K. E. and Gordon, D. C.: 1998, UV-B and Canada's Forests: State of Science and Risk. Pp 75-85 in Legge, A. H. and L.L. Jones (ed.), *Proceedings Air and Waste Management Association Speciality Conference on Emerging Air Issues for the 21st Century*, Calgary
- Percy, K. E., Jagels, R., Marden, S., McLaughlin, C. K., Carlisle, J.: 1993, *Can. J. For. Res.* **23**, 1472.
- Persson, H. and Majdi, H.: 1995, *Water, Air, and Soil Pollut.* **85**, 1287.
- Peterson, D. L., Silsbee, D. G., Poth, M., Arbaugh, J., Biles, F. E.: 1995, *J. Air Waste Manage. Assoc.* **45**, 36.
- Peterson, D. L., Arbaugh, M. J., Robinson, L. J.: 1991, *Holoc.* **1**, 50.
- Pooviah, B. W.: 1988, *Senescence and Aging in Plants*, Academic Press, Inc.

- Powell, D. S., Faulkner, J. L., Darr, D. R., Zhu, Z., MacCleery, D. W.: 1992, *Forest Resources of the United States*. USDA Forest Service Gen. Tech. Report RM-234, 132 pp.
- Prinz, B.: 1987, *Environment* 29, 10.
- Pronos, J., Vogler, D. R., Smith, R. S.: 1978, An evaluation of ozone injury to pines in the southern Sierra Nevada. USDA Forest Service Forest Pest Management Report 78-1, San Francisco.
- Rehfuess, K. E.: 1981, *Forstwiss. Centralbl.* 100, 363.
- Reiners, W. A. and Lang, G. E.: 1979, *Ecol.* 60, 403.
- Rennenberg, H., Kreutzer, K., Papen, H., Weber, P.: 1998, *New Phytol.* 139, 71.
- Reuss, J. O. and Johnson, D. W.: 1986, *Acid Deposition and the Acidification of Streams and Waters*. Springer-Verlag.
- Richardson, C. T., Sasek, T. W., Fendick, E. A.: 1992, *Environ. Toxicol. Chem.* 11, 1105.
- Richter, D. D. and Markewitz, D.: 1995, pp. 315-336 in S. Fox and R. A. Mickler (eds.), *Impacts of Air Pollutants on Southern Pine*. Springer-Verlag.
- Richter, D. D., Markewitz, D., Wells, C. G., Allen, H. L., April, R., Heine, P. R., Urrego, B.: 1994, *Ecology* 75, 1463.
- RMCC: 1990, *The 1990 Canadian long-range transport of air pollutants and acid deposition assessment report: Part I Executive Summary*. Federal-Provincial Research and Monitoring Coordinating Committee, Ottawa.
- Robarge, W. P., Pye, J. M., Bruck, R. I.: 1989, *Plant and Soil* 114, 19.
- Robarge, W. P. and Johnson, D. W.: 1992, *Advances in Agronomy* 47, 1.
- Roberts, D. M. and Harmon, A. C.: 1992, *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 43, 375.
- Robitaille, G., Boutin, R., Lachance, D.: 1995, *Can. J. For. Res.*, 25, 577.
- Roy, G., Robitaille, G., Gagnon, G.: 1985, *Phytoprotection* 66, 91.
- Royle, D. D. and Lathrop R. G.: 1997, *For. Sci.* 43, 327.
- Samuelson, L. J. and Kelly, J. M.: 1997, *New Phytologist* 136, 255.
- Sasek, T. W. and Flagler, R. B.: 1995, *Impacts of Air Pollutants on Southern Pine Forests*. Springer-Verlag. pp. 424-466.
- Sasek, T. W., Richardson, C. J., Fendick, E. A., Bevington, S. R., Kress, L. W.: 1991, *For. Sci.* 37, 1078.
- Saxena, V. K., and Lin, N. H.: 1990, *Atmos. Envir.* 24A, 329.
- Schaberg, P. G., Perkins, T. D., McNulty, S. G.: 1997, *Can. J. For. Res.* 27, 1622.
- Schemenauer, R. S.: 1986, *Atmos. Ocean* 24, 303.
- Scherbatskoy, T. de W.: 1989, *Ionic Relations of Leaf Cuticles*, Ph.D. Thesis, University of Vermont.
- Schroeder, W. H., Bauch, J., Endeward, R.: 1988, *Trees* 2, 96.
- Schulze, E. D.: 1989, *Science* 244, 776.
- Schulze, E. D., De Vries, W. Hauhs, M., Rosen, K., Rasmussen, L., Tamm, C. O., Nilsson J.: 1989, *Water, Air, and Soil Pollut.* 48, 451.
- Schulze, E. D., Robichaux, R. H., Grace, J., Rundel, P. W., Ehleringer, J. R.: 1987, *BioScience* 37, 30.
- Schutt, P. and Cowling, E. B.: 1985, *The Amer. Phytopathological Soc.* 69,
- SCOPE: 1992, *Effects of increased ultraviolet radiation on biological systems*. Scientific Committee on Problems in the Environment, Paris.
- Sheffield, R. M. and Cost, N. D.: 1987, *J. For.* 87, 29.

- Sheffield, R. M. et al.: 1985, USDA For. Serv. Res. Bull. SE-83.
- Shepard, M. R., Lee, C. E., Woosley, R. S., Butcher, D. J.: 1995, *Microchemical J.* **52**, 118.
- Shortle, W. C. and Bondietti, E. A.: 1992, *Water, Air, and Soil Pollut.* **61**, 253.
- Shortle, W. C., Smith, K. T., Minocha, R., Alexeyev, V. A.: 1995, *J. Biogeography* **22**, 467.
- Shortle, W. C. and Smith, K. T.: 1988, *Science* **240**, 1017.
- Shriner, D. S., Heck, W. W., McLaughlin, S. B., Johnson, D. W., Peterson, C. E.: 1990, *NAPAP State of Science and Technology*, National Acidic Precipitation Assessment Program.
- Skarby, L., Troeng, E., Bostrom, C. A.: 1987, *For. Sci.* **33**, 801.
- Skarby, L., Ro-Poulsen, H., Wellburn, F.A.M., Sheppard, L. J.: 1998, *New Phytol.* **139**, 109.
- Smith, W., and Mangold, R.: 1996, *Estimating status and change in forest health*, U.S. Dep. Agric. Forest Service Fact Sheet.
- Smith, G. F., and Nicholas, N. S.: 1999, *Castanea* (in press).
- Stark, R. W.: 1965, *Annu. Rev. Entomol.* **10**, 303.
- Sterba, H.: 1996, *Forest decline and growth trends in Central Europe - a review*. Berlin: Springer-Verlag.
- Stienen, H. and Bauch, J.: 1988, *Plant and Soil* **106**, 231.
- Stohlgren, T. J. and Parsons, D. J.: 1987, *Atmos. Environ.* **21**, 1969.
- Stow, T. K., Allen, H. L., Kress, L. W.: 1992, *For. Sci.* **38**, 102.
- Stoyenhoff, J., Witter, J., Leutscher, B.: 1998, *Forest Health in the New England States and New York*, University of Michigan, 31 pp.
- Swank, W. T. and Vose, J. M.: 1997, *Global Biogeochemical Cycles* **11**, 657.
- Taylor, G. E.: 1994, *J. Environ. Qual.* **23**, 63.
- Teskey, R. O.: 1995, pp. 467-490 in S. Fox and R. A. Mickler (eds.), *Impacts of Air Pollutants on Southern Pine*. Springer-Verlag.
- Teskey, R. O., Dougherty, P. M., Mickler, R. A.: 1998, pp. 279-290 in R. A. Mickler and S. Fox (eds.), *The Productivity and Sustainability of Southern Forest Ecosystems in a Changing Environment*. Springer-Verlag.
- Thomson, D. J.: 1995, *Science* **268**, 59.
- Timmel, T.E. 1996. pp 1907-1969 in *Compression Wood Vol 3*. Springer-Verlag.
- Tingey, D. T. and Hogsett, W. E.: 1985, *Plant Physiol.* **77**, 944.
- Trewavas, A. J. and Malho, R.: 1997, *The Plant Cell* **9**, 1181.
- Ulrich, B.: 1984, *Atmos. Env.* **18**, 621.
- Ulrich, B. and Matzner, E.: 1986, *Experientia* **42**, 344.
- USDA.: 1997, *Forest Insect and Disease Conditions in the United States 1996*. Forest Service Health Production, Washington, D.C., 87 p.
- USDA Forest Service: 1996, *Sampling and plot design*, Forest health monitoring fact sheet 1909.14, USDA Forest Service, Washington, DC.
- Vann, D. R., Strimback, G. R., Johnson, A. H.: 1992, *For. Ecol. Manag.* **51(1-3)**, 69.
- Wallin, G. and Skarby, L.: 1992, *Trees* **6**, 128.
- Wargo, P. M.: 1972, *Phytopathology* **62**, 1278.
- Waring, G. L. and Cobb, N. S.: 1992, *Insect-Plant Interactions*. pp. 167-226.
- Waring, R. H.: 1987, *BioScience* **37**, 569.
- Waring, R. H. and Pitman: 1985, *Ecol.* **66**, 889.
- Watmough, S. A.: 1997, *Envir. Rev.* **5**, 181.

- Watson, R. T., Zinyowera, M. C., Moss, R. H., Dokken, D. J.: 1998, *The Regional Impacts of Climate Change, An Assessment of Vulnerability*. Cambridge University Press.
- Weeks, M. W., and Smoronok, D.: 1993, *Aerial assessment of red spruce and balsam fir conditions in the Adirondack region of New York, etc.* USDA-USFS Technical Report NA-TP-16-93, 141 p.
- Wesselink, L. G., Meiwes, K. J., Matzner, E., Stein, A.: 1995, *Environ. Sci. Technol.* **29**, 51.
- Wessman, C. A., Aber, J. D., Peterson, D. L.: 1989, *Int. J. Remote Sensing* **10**, 1293.
- Wisniewski, J.: 1982, *Water, Air, Soil Pollut.*, **17**, 361.
- Witkowski, Z., Madziara-Borusiewicz, K., Plonka, P., Zurek, Z.: 1987, *Ekologia Polska* **35**, 465.
- Wulschleger, S. D., Norby, R. J., Gunderson C. A.: 1997, pp. 79-100 in L. H. Allen, *Advances in Carbon Dioxide Effects Research*. ASA Special Publication No. 61, X.
- Foster, N. W. and Arp, P. A.: 1993, *Can. J. For. Res.*, **23**, 617.
- Yin, X., Foster, N. W., Morrison, I. K., Arp, P.A.: 1994, *Can. J. For. Res.*, **24**, 1567.
- Yoder, B. J., Ryan, M. G., Waring, R. H., Schoettle, A. W., Kaufmann, M. R.: 1994, *For. Sci.* **40**, 513.
- Zahner, R., Saucier, J. R. and Myers, R. K.: 1989, *Can. J. For. Res.* **19**:612.
- Zhao, X. J., Sucoff, E., Stadelmann, E. J.: 1987, *Plant Physiol.* **83**, 159.
- Zoetl, H. W. and Huettl, R. F.: 1986, *Water, Air, and Soil Pollut.* **31**, 449.

Acknowledgements

The authors wish to express their sincere appreciation to Dr. R. Vet, Atmospheric Environment Service, Environment Canada, Downsview, Ontario and to Dr. T. Dann, Environmental Protection Service, Environment Canada, Ottawa, Ontario for calculation and mapping of SO₄, NO₃, H ion depositions and cumulative O₃ sums. The senior author thanks the U.S. Forest Service for support of some aspects of his basic research reported herein and the Department of Energy, Bioenergy Feedstock Development Program at Oak Ridge National Laboratory for partial support of synthesis activities. This manuscript is Publication No. 4892 of the Environmental Sciences Division, Oak Ridge National Laboratory. ORNL is operated by Lockheed Martin Energy Research for the U.S. Department of Energy under contract No. DE-ACO5-96OR22464.